

Symptoms of Dissociation in Humans Experiencing Acute, Uncontrollable Stress: A Prospective Investigation

Charles A. Morgan III, M.D., M.A.

Maj. Gary Hazlett, Psy.D.

Sheila Wang, Ph.D.

E. Greer Richardson, Jr., M.D.

Paula Schnurr, Ph.D.

Steven M. Southwick, M.D.

Objective: Peritraumatic dissociation has been associated with subsequent development of posttraumatic stress disorder, but supporting data have been largely retrospective. The current study was designed to assess the nature and prevalence of dissociative symptoms in healthy humans experiencing acute, uncontrollable stress during U.S. Army survival training.

Method: In study 1, 94 subjects completed the Clinician-Administered Dissociative States Scale after exposure to the stress of survival training. In study 2, 59 subjects completed the Brief Trauma Questionnaire before acute stress and the dissociative states scale before and after acute stress. A randomly selected group of subjects in study 2 completed a health problems questionnaire after acute stress.

Results: In study 1, 96% of subjects reported dissociative symptoms in response to acute stress. Total scores, as well as individual item scores, on the dissociation

scale were significantly lower in Special Forces soldiers compared to general infantry troops. In study 2, 42% of subjects reported dissociative symptoms before stress and 96% reported them after acute stress. Dissociative symptoms before and after stress were significantly higher in individuals who reported a perceived threat to life in the past. Forty-one percent of the variance in reported health problems was accounted for by poststress dissociation scores.

Discussion: Symptoms of dissociation were prevalent in healthy subjects exposed to high stress. Stress-hardy individuals (Special Forces soldiers) experienced fewer symptoms of dissociation, compared to individuals who were less hardy. These data support the idea that the nature of response to previously experienced threatening events significantly determines the nature of psychological and somatic response to subsequent stress.

(*Am J Psychiatry* 2001; 158:1239–1247)

Although many people are exposed to trauma, only some individuals develop posttraumatic stress disorder (PTSD); most do not (1–4). Further, in a large proportion of individuals who do develop the disorder, the disorder fully remits over time. These findings suggest that chronic PTSD may represent a specific type of adaptation to trauma that is not typical of most individuals, but rather is seen in individuals who have a particular kind of vulnerability to stress. Researchers have identified several specific psychosocial risk factors for PTSD, including a history of exposure to traumatic events, exposure to multiple traumatic events, exposure to childhood sexual or physical trauma, and the subjective experience of fear for one's life (5–7). However, because these factors still leave a great deal of the variance unexplained, investigations evaluating risk factors for the development of PTSD continue.

Over the past decade, numerous investigators have begun to reexamine the relationship between trauma, symptoms of dissociation, and the development of PTSD (8–17). Taken together, the data from several studies suggest that peritraumatic symptoms of dissociation (i.e., symptoms of dissociation experienced during and for a short time immediately after exposure to a traumatic event) represent a significant risk factor for the subsequent development of

PTSD. Symptoms of dissociation also appear to be significantly related to somatic complaints reported by victims of trauma (18) and may play a significant role in the reported relationship between PTSD and physical health (19).

Although intriguing, these findings cannot be interpreted with a high degree of certainty because they were based on retrospective accounts. Consequently, it remains unclear whether: 1) dissociation serves as a risk factor for subsequent development of PTSD, 2) trauma independently causes dissociation and PTSD, or 3) individuals with current dissociative symptoms are biased toward overreporting the presence and severity of dissociative symptoms at the time of a past trauma.

Although peritraumatic dissociation appears to be a risk factor for stress-related illness, surprisingly little is known about its nature and frequency in individuals (such as military personnel) who are at high risk for exposure to trauma and PTSD. At present, it is not known whether the propensity to dissociate is the result of trauma exposure or whether it represents a trait that not only predisposes an individual to dissociate during stress but also renders that person more vulnerable to the development of PTSD. Examining individuals' propensity to dissociate before, and in response to, stress exposure—before the development

of PTSD—would provide data about whether dissociation is a relatively common response to realistic threats and whether there is a relationship between history of trauma exposure and propensity to dissociate when confronted with high stress. This type of information would help clarify the nature of the relationship between dissociation and PTSD.

This study was part of a larger investigation examining the neuroendocrine profiles of healthy soldiers participating in survival training conducted by the U.S. Army (20, 21). The design permitted an assessment of the presence and severity of state dissociative symptoms in healthy military subjects before and after exposure to highly intense stress. The design also permitted an assessment of whether factors such as the amount (or type) of previous trauma, as well as the nature of response to previously experienced trauma, would significantly contribute to symptoms of dissociation before and during acute stress.

The U.S. Army's survival training course is among the most difficult and rigorous training programs in the U.S. Armed Forces. Several factors make the survival course an ideal environment in which to study the relationship between highly intense stress and symptoms of dissociation. First, the course participants represent a healthy, nonclinical group of subjects who are at high risk for exposure to military-related trauma and, consequently, at risk for the development of combat-related stress disorders such as PTSD. Second, the course allows for a highly controlled and uniform application of stress across subjects. Third, the training scenario represents the best analog of realistic military stress, as it is multidimensional (combining psychological, physical, and environmental stress) and results in neurobiological alterations that are on a par with those that occur in actual threat-to-life experiences (see the last paragraph in this section) (20, 21).

The survival course is 19 days long and includes a low-stress didactic (classroom) phase and a highly stressful experiential phase. During the experiential phase, students are confronted with a variety of stressors, including semistarvation, sleep deprivation, lack of control over personal hygiene, and external control over movement, social contact, and communication. The experiential phase provides students with the opportunity to apply their training in a controlled, albeit stress-laden, environment.

Previous investigations have provided robust evidence that the stress of the experiential phase of the survival school training produces significant alterations in neuroendocrine responses, including changes in glucocorticoids, catecholamines, gonadal steroids, and neuropeptide Y (20, 21). The magnitude of these responses is analogous to that produced in real-world life-threatening situations and is also similar to that found in previous studies of high-stress military training (nocturnal landings on an aircraft carrier, military free-fall, and Ranger training). As such, survival school training provides an ethologically realistic setting for studying neurobio-

logical aspects of acute stress in humans and is well suited for a prospective examination of the relationship between acute stress and psychological symptoms of dissociation.

Method

Subjects and Procedure

Two studies with separate groups of subjects were conducted. The subjects of study 1 were 50 Special Forces and 44 general infantry soldiers (mean age=31.6 years, SD=4.8, and mean age=28.9 years, SD=5.1, respectively). The subjects of study 2 were a separate group of 18 Special Forces and 41 general infantry soldiers (mean age=27.1 years, SD=4.4, and mean age=22.0, SD=3.6, respectively). Special Forces soldiers—also known as Green Berets—are individuals who have successfully completed the Special Forces Selection and Assessment course, followed by the Special Forces Qualification Course. These training hurdles involve high levels of challenge to physical endurance, academic abilities, and stress tolerance. Only 25% of applicants are ultimately successful in the Special Forces Selection and Assessment course. Thus, Special Forces soldiers represent a highly selected population of military subjects.

Per course requirements, all subjects received medical and psychological clearance before in-processing to survival school at Fort Bragg, N.C. All were free of illicit substances. After completing in-processing to the survival school, subjects were given a description of the nature and purpose of the study by the principal investigator (C.A.M.). Each gave written, informed consent to participate in the study. Each understood that refusal to participate in the study would not adversely affect their standing in the survival course. Sixteen subjects declined to participate in the study. Eleven of the 16 stated that they did not enroll in the study because they were not confident that research information would remain separate from their military records. They worried that such information, if inadvertently included in their military records, might jeopardize their subsequent evaluations. The remaining five did not offer an explanation for refusing to participate in the study.

Subjects in study 1 were administered the Clinician-Administered Dissociative States Scale after completing the experiential phase of the survival training. The Clinician-Administered Dissociative States Scale is a reliable, valid, self-report instrument designed to assess state symptoms of dissociation in response to a specified stressor (22). The scale includes 19 self-report items and eight observer-rated (clinician-rated) items. Subjects rated the intensity of the dissociative experiences on a scale from 0, not present, to 4, extreme. The scores for individual items were summed to produce the total score. Subjects were instructed to respond to the scale items by using the experiential period of the survival training as the reference point. Due to the design and nature of the survival training, it was not possible to accurately determine ratings for the eight observer-rated items. Thus, only the subjective items were rated. This modification is compatible with other self-report inventories and does not alter the validity or reliability of the Clinician-Administered Dissociative States Scale (22; J.D. Bremner, personal communication, 1999).

Subjects in study 2, a separate cohort from those in study 1, completed the Clinician-Administered Dissociative States Scale and the Brief Trauma Questionnaire during the week of classroom activities before exposure to acute stress. These subjects completed the dissociation scale again at the conclusion of the experiential phase of the training. When filling out the dissociation scale at baseline, subjects were instructed to use the two previous days of classroom activities as their reference point. When

TABLE 1. Severity of Individual State Dissociative Symptoms After Uncontrollable Stress^a in Healthy Special Forces Soldiers and General Infantry Soldiers (Study 1)

	Soldiers Reporting Presence of Symptom		Symptom Severity				Analysis ^c		
			Special Forces Soldiers (N=50)		General Infantry Soldiers (N=44)				
	N	%	Mean	SD	Mean	SD	t	df	p
Clinician-Administered Dissociative States Scale ^b									
Scale items measuring dissociative symptoms									
1. Things seemed to move in slow motion.	54	57	0.9	1.0	1.2	1.2	1.7	92	n.s.
2. Things seemed unreal, as if in a dream.	49	52	0.7	0.9	1.0	1.1	1.2	92	n.s.
3. You had a feeling of separation from what was happening, as if you were watching a movie or a play or as if you were an automaton.	39	41	0.6	1.0	1.1	1.2	2.3	92	<0.02
4. You felt as if you were watching things from outside your body.	21	22	0.1	0.4	0.6	0.9	3.1	56	<0.003
5. You felt as if you were watching the situation as an observer or a spectator.	28	30	0.3	0.5	0.7	1.0	2.4	65	<0.01
6. You felt disconnected from your body.	21	22	0.2	0.6	0.7	1.0	2.6	73	<0.01
7. Your sense of your own body seemed changed.	37	39	0.5	0.8	1.1	1.3	2.8	74	<0.007
8. People seemed motionless, dead, or mechanical.	41	44	0.6	0.9	1.1	1.2	2.0	78	<0.05
9. Objects looked different than you would have expected.	41	44	0.4	0.6	1.3	1.2	4.7	64	<0.0001
10. Colors seemed diminished in intensity.	48	51	0.7	1.0	1.3	1.2	2.4	92	<0.02
11. You saw things as if in a tunnel or through a wide-angle photographic lens.	32	34	0.4	0.7	1.1	1.2	3.2	70	<0.002
12. Things seemed to take much longer than you would have expected.	62	66	1.5	1.3	1.7	1.6	0.5	92	n.s.
13. Things seemed to happen very quickly, as if there were a lifetime in a moment.	43	46	1.0	1.3	1.1	1.4	0.4	92	n.s.
14. Things happened that you were unable to account for later.	54	57	0.8	0.8	1.3	1.2	2.3	79	<0.02
15. You spaced out or lost track of what was going on.	61	65	0.9	1.1	1.8	1.2	3.8	92	<0.0002
16. Sounds almost disappeared or became stronger than you would have expected.	50	53	0.8	1.1	1.6	1.4	3.2	92	<0.002
17. Things seemed very real, as if there were a special sense of clarity.	57	61	0.9	1.2	1.9	1.3	4.3	92	<0.0001
18. You felt as if you were looking at the world through a fog; people and objects appeared far away or unclear.	92	98	0.2	0.5	1.0	1.3	4.0	57	<0.0001
19. Colors seemed much brighter than you would have expected.	28	30	0.2	0.5	0.9	1.2	3.8	63	<0.0006
Scale total			11.6	9.6	22.5	16.4	3.8	67	<0.0001

^a Uncontrollable stress presented by the experiential phase of survival training course, during which soldiers were subject to semi-starvation, sleep deprivation, lack of control over personal hygiene, and external control over movement, social contact, and communication.

^b Intensity of dissociative experiences rated on a scale from 0, not present, to 4, extreme. Scores for individual items were summed to produce the total score.

^c When indicated degrees of freedom (df) differed from 92, the Levine test was rejected and equal variances in the groups were not assumed.

filling out the poststress assessment, subjects were given instructions to complete the scale in response to the two previous days of the experiential phase.

Subjects in study 2 also completed the Brief Trauma Questionnaire in the didactic phase of the course, prior to stress exposure. The Brief Trauma Questionnaire is a valid, reliable, self-report instrument designed to assess the number and types of traumatic events to which an individual has been exposed in life (23). The scale also permits an assessment of whether an individual suffered "physical injury" or experienced "fear for one's life" in relation to a previous trauma. Endorsement of either of these items is interpreted as exposure to a traumatic event, as outlined in criterion A of the DSM-IV criteria for posttraumatic stress disorder.

Finally, just after completing the experiential phase of the survival course, a randomly selected group of subjects in study 2 (N=36) were administered the Health Problem List. The Health Problem List is a valid and reliable self-report instrument comprising 80 items designed to assess current somatic symptoms (such as heartburn, chest pain, ringing in ears, etc.) (24). The to-

tal score is the number of items endorsed. In this paper this score will be referred to as the somatic symptom score to avoid any misconception that this instrument reflects a medical history inventory.

Data Analysis

In study 1, t tests were used to compare Special Forces and general infantry soldiers on poststress Clinician-Administered Dissociative States Scale scores. In study 2, paired t tests were used to examine change over time in dissociation scale item scores and total score. In addition, independent t tests were used to evaluate whether dissociation scale item scores differed between Special Forces and general infantry soldiers. Pearson and point-biserial correlations were used for univariate comparisons within each time point (prestress and poststress) in order to select variables for multivariate analysis. Simultaneous multiple hierarchical regression analyses were used to examine predictors of Clinician-Administered Dissociative States Scale scores after the stress of the experiential phase. A p value of 0.05 was used for interpreting statistical significance.

TABLE 2. Severity of Individual State Dissociative Symptoms Before and After Uncontrollable Stress^a in Healthy Special Forces Soldiers and General Infantry Soldiers (Study 2)

Clinician-Administered Dissociative States Scale ^b	Soldiers Reporting Presence of Symptom		Symptom Severity				Analysis ^c		
	N	%	Special Forces Soldiers (N=18)		General Infantry Soldiers (N=41)		t	df	p
			Mean	SD	Mean	SD			
Scale items measuring dissociative symptoms									
1. Things seemed to move in slow motion.									
Before stress	16	27	0.3	0.7	0.5	1.0	0.6	57	n.s.
After stress	36	61	0.8	1.1	1.2	1.1	1.3	57	n.s.
2. Things seemed unreal, as if in a dream.									
Before stress	14	24	0.7	0.9	1.0	1.1	1.6	56	n.s.
After stress	37	63	0.6	0.8	1.2	1.2	2.2	57	<0.03
3. You had a feeling of separation from what was happening, as if you were watching a movie or a play or as if you were an automaton.									
Before stress	11	19	0.1	0.2	0.4	0.9	2.3	50	<0.03
After stress	30	51	0.4	0.6	1.0	1.1	2.1	57	<0.03
4. You felt as if you were watching things from outside your body.									
Before stress	9	15	0.0	0.0	0.3	0.8	2.7	40	<0.01
After stress	14	24	0.1	0.2	0.4	0.8	2.1	57	n.s.
5. You felt as if you were watching the situation as an observer or a spectator.									
Before stress	11	19	0.3	0.5	0.7	1.1	2.9	57	<0.01
After stress	19	32	0.1	0.2	0.7	1.0	2.6	57	<0.01
6. You felt disconnected from your body.									
Before stress	9	15	0.1	0.2	0.2	0.6	1.5	56	n.s.
After stress	25	42	0.2	0.5	0.8	0.9	2.7	57	<0.008
7. Your sense of your own body seemed changed.									
Before stress	14	24	0.2	0.4	0.4	0.8	1.3	56	n.s.
After stress	31	53	0.5	0.8	1.4	1.4	2.6	57	<0.01
8. People seemed motionless, dead, or mechanical.									
Before stress	8	14	0.0	0.0	0.3	0.8	2.1	40	<0.05
After stress	28	47	0.3	0.7	1.4	1.4	2.9	57	<0.005
9. Objects looked different than you would have expected.									
Before stress	14	24	0.2	0.5	0.4	0.9	0.9	56	n.s.
After stress	28	47	0.4	0.7	1.1	1.3	2.2	55	<0.04
10. Colors seemed diminished in intensity.									
Before stress	11	19	0.1	0.2	0.3	0.7	1.9	54	n.s.
After stress	22	37	0.3	0.6	0.7	1.1	1.8	55	n.s.
11. You saw things as if in a tunnel or through a wide-angle photographic lens.									
Before stress	11	19	0.1	0.5	0.4	0.9	1.4	55	n.s.
After stress	25	42	0.2	0.5	1.0	1.2	2.7	57	<0.009
12. Things seemed to take much longer than you would have expected.									
Before stress	14	24	0.4	0.8	0.4	0.9	0.9	57	n.s.
After stress	54	92	2.5	1.0	2.4	1.3	0.1	57	n.s.
13. Things seemed to happen very quickly, as if there was a lifetime in a moment.									
Before stress	11	19	0.1	0.3	0.3	0.8	1.4	56	n.s.
After stress	21	36	0.3	1.0	0.8	1.2	1.6	40	n.s.
14. Things happened that you were unable to account for later.									
Before stress	11	19	0.0	0.0	0.5	1.1	1.8	57	n.s.
After stress	35	59	0.8	0.9	1.0	1.1	0.8	57	n.s.
15. You spaced out or lost track of what was going on.									
Before stress	18	31	0.2	0.4	0.6	1.1	2.0	56	n.s.
After stress	41	69	1.0	0.9	1.5	1.3	1.6	47	n.s.
16. Sounds almost disappeared or became stronger than you would have expected.									
Before stress	11	19	0.1	0.2	0.4	0.9	2.1	51	<0.03
After stress	32	54	0.7	1.0	1.1	1.2	1.3	57	n.s.
17. Things seemed very real, as if there were a special sense of clarity.									
Before stress	14	24	0.1	0.2	0.4	0.9	2.3	50	<0.02
After stress	39	66	1.1	1.2	1.1	1.1	0.1	57	n.s.

(continued)

TABLE 2. Severity of Individual State Dissociative Symptoms Before and After Uncontrollable Stress^a in Healthy Special Forces Soldiers and General Infantry Soldiers (Study 2) (continued)

Clinician-Administered Dissociative States Scale ^b	Soldiers Reporting Presence of Symptom		Symptom Severity				Analysis ^c		
	N	%	Special Forces Soldiers (N=18)		General Infantry Soldiers (N=41)		t	df	p
			Mean	SD	Mean	SD			
18. You felt as if you were looking at the world through a fog; people and objects appeared far away or unclear.									
Before stress	8	14	0.0	0.0	0.2	0.7	2.2	40	<0.03
After stress	23	39	0.3	0.8	0.7	0.9	1.9	57	n.s.
19. Colors seemed much brighter than you would have expected.									
Before stress	11	19	0.0	0.0	0.4	0.9	2.6	57	<0.01
After stress	17	29	0.3	0.8	0.4	0.8	0.7	57	n.s.
Scale total									
Before stress			1.6	2.5	6.0	11.1	1.7	58	n.s.
After stress			9.7	5.8	21.3	14.6	2.4	56	<0.003

^a Uncontrollable stress presented by the experiential phase of survival training course, during which soldiers were subject to semi-starvation, sleep deprivation, lack of control over personal hygiene, and external control over movement, social contact, and communication.

^b Intensity of dissociative experiences rated on a scale from 0, not present, to 4, extreme. Scores for individual items were summed to produce the total score.

^c When indicated degrees of freedom (df) differed from 57, the Levine test was rejected and equal variances in the groups were not assumed.

Results

Study 1

Total mean Clinician-Administered Dissociative States Scale scores after stress were significantly lower in the Special Forces subjects, compared to the general infantry subjects. As shown in Table 1, comparisons between the two groups for each dissociation scale item revealed significant differences on all but four items (items 1, 2, 12, and 13). Table 1 also shows that the most common symptoms of dissociation for the overall group of subjects were those in items 18, 12, and 15 ("looking at the world through a fog" [98%], the experience taking longer than expected [66%], and "spacing out" [65%], respectively).

Study 2

There was a substantial overall increase in reports of dissociative symptoms from before to after the acute stress of the experiential phase, indicating that the experiential phase of the survival course was highly stressful (Table 2). Pre- and poststress scores were significantly correlated ($r=0.45$, $df=57$, $p<0.001$). Paired t tests (not shown in Table 2) indicated that significant within-subject differences existed between the prestress and the poststress total scores on the Clinician-Administered Dissociative States Scale ($t=6.8$, $df=56$, $p<0.0001$). Significant within-subject increases in scores were also observed on the individual scale items, with the exception of items 4 and 19 (item 1: $t=4.1$, $df=56$, $p<0.0001$; item 2: $t=4.2$, $df=56$, $p<0.0001$; item 3: $t=3$, $df=56$, $p<0.003$; item 4: $t=0.7$, $df=56$, $p<0.40$; item 5: $t=2.3$, $df=56$, $p<0.03$; item 6: $t=3.8$, $df=56$, $p<0.0001$; item 7: $t=4.6$, $df=56$, $p<0.0001$; item 8: $t=5.0$, $df=56$, $p<0.0001$; item 9: $t=3.1$, $df=56$, $p<0.003$; item 10: $t=2.7$, $df=56$, $p<0.01$; item 11: $t=2.8$, $df=56$, $p<0.008$; item 12: $t=12.4$, $df=56$, $p<0.00001$; item 13: $t=2.4$, $df=56$, $p<0.02$; item 14: $t=3.5$, $df=56$, $p<0.001$; item 15: $t=5.0$, $df=56$, $p<0.0001$;

item 16: $t=4.2$, $df=56$, $p<0.0001$; item 17: $t=4.7$, $df=56$, $p<0.002$; item 18: $t=3.2$, $df=56$, $p<0.002$; and item 19: $t=0.8$, $df=56$, $p<0.40$).

Table 2 indicates the most common symptoms of dissociation before and after stress exposure for the group as a whole and also compares the Special Forces and general infantry groups on each of the pre- and poststress dissociation scale items. As in study 1, alterations in perception of time, perception of the environment, and clarity of thought (for example, items 12, 15, and 17) were among the ten most common symptoms reported by subjects. Table 2 also shows the results of the independent t tests comparing the two groups' scores on each of the dissociation scale items before and after stress. As in study 1, scores on items 3, 5–9, and 11 differed significantly between the Special Forces and the general infantry soldiers after stress exposure, whereas items 12 and 13 did not. Unlike the findings in study 1, the study 2 findings showed that the two groups did not differ significantly in their poststress scores on items 4 and 14–19. The probability values for these items were at the 0.09 significance level, and the findings may be the result of a lack of power given the smaller number of Special Forces soldiers in study 2.

Table 3 shows the frequency of and types of trauma that subjects in study 2 had previously experienced. The subjects provided this information before beginning the experiential phase of the survival course. More than 90% of the subjects reported previous exposure to a potentially traumatic stressor.

Table 4 presents the results of univariate tests of association for baseline and poststress Clinician-Administered Dissociative States Scale scores with Special Forces membership and trauma exposure. To facilitate comparisons among the predictors, the data are presented as correlations. All results are point-biserial correlations except for those involving number of traumas, which are Pearson

TABLE 3. Traumas Ever Experienced by 59 Healthy Special Forces and General Infantry Soldiers in a Study of Reaction to Uncontrollable Stress^a (Study 2)

Trauma Ever Experienced ^b	Trauma Exposure			
	Yes		No	
	N	%	N	%
Traumatic event				
Exposure to any trauma	55	93	4	7
War	15	25	44	75
Accident	23	39	36	61
Disaster	22	37	37	63
Illness	1	2	58	98
Early trauma	13	22	46	78
Mugging	22	37	37	63
Sexual abuse	5	8	54	92
Fear-provoking situation	28	47	31	53
Witnessing trauma	37	63	22	37
Experiencing "fear for life, horror, helplessness"	29	49	30	51
Trauma that resulted in physical injury	9	15	50	85
Number of traumatic events				
None	4	7		
One	13	22		
Two	5	8		
Three	15	25		
Four	5	8		
Five	11	19		
Six	4	7		
Seven	2	3		

^a Uncontrollable stress presented by the experiential phase of survival training course, during which soldiers were subject to semi-starvation, sleep deprivation, lack of control over personal hygiene, and external control over movement, social contact, and communication.

^b Trauma experience measured with the Brief Trauma Questionnaire, completed before subjects began the experiential phase of survival training course.

correlations. Only two variables were correlated with dissociation scale scores at one or both time points. Special Forces membership was unrelated to the prestress score but was related to the poststress score, replicating the findings of study 1. Perceived life threat during a trauma was related to higher dissociation scale scores both before and after stress.

The fact that only two dichotomous variables were related to dissociation scale scores provided an opportunity to examine change in scores after stress exposure in subtypes of individuals who differed in potential vulnerability. Three dummy variables were created to reflect the following natural groups: Special Forces soldiers with previous exposure to a perceived threat to life (N=8), general infantry soldiers with no prior perceived threat to life (N=20), and general infantry soldiers with prior exposure to a perceived threat to life (N=20). The Special Forces subjects with no prior perceived threat to life (N=9) were selected as the reference group because we assumed they would have the least reaction to the stress of the experiential phase of the survival course. The three dummy variables were used in a multiple regression analysis predicting poststress Clinician-Administered Dissociative States Scale scores, while controlling for prestress scores. This approach is analogous to predicting change in dissociation scale scores from before to after stress exposure. Rela-

TABLE 4. Correlations Between Predictors and Severity of State Dissociative Symptoms Before and After Uncontrollable Stress^a in Healthy Special Forces and General Infantry Soldiers (Study 2)

Predictor	Correlation ^b With Severity of State Dissociative Symptoms, Rated With the Clinician-Administered Dissociative States Scale	
	Before Stress	After Stress
Special Forces membership	-0.20	-0.38*
Previous exposure to perceived threat to life	0.38*	0.36*
Injury during previous trauma	0.14	-0.10
Childhood physical abuse	0.07	-0.01
Number of lifetime traumas	0.06	0.12

^a Uncontrollable stress presented by the experiential phase of survival training course, during which soldiers were subject to semi-starvation, sleep deprivation, lack of control over personal hygiene, and external control over movement, social contact, and communication.

^b Point-biserial correlations for all predictors except number of traumas, for which Pearson correlations were used.

*p<0.01.

tive to the Special Forces subjects without a prior threat to life, only the general infantry subjects who had experienced a threat to life had a significantly greater score after stress exposure. The unstandardized regression coefficients, which represent the difference in adjusted mean change in scores between each group and the group of Special Forces subjects with no prior threat to life, were as follows: Special Forces soldiers with a prior threat to life (B=-1.48, n.s.), general infantry soldiers with no prior threat to life (B=3.46, n.s.), and general infantry soldiers with a prior threat to life (B=12.97, p<0.005). To facilitate interpretation of the regression analysis, prestress and poststress Clinician-Administered Dissociative States Scale scores are represented in Figure 1.

Scores on several dissociation scale items, however, were significantly different between groups before acute stress. These items pertained to a sense of separation from one's body—watching things as if outside one's body—and to altered perception of other individuals and the environment. Nevertheless, scores on many items (items 3 and 5-11) were once again found to be significantly lower in Special Forces subjects than in general infantry subjects.

No significant correlations were observed between age, number of traumas, the baseline Clinician-Administered Dissociative States Scale score, and the somatic symptoms score. Significant, positive correlations were observed between the somatic symptoms score and the poststress dissociation scale score (r=0.54, df=34, p<0.02) as well as between the somatic symptoms score and the change in the dissociation scale score from the baseline to the poststress assessment (r=0.67, df=34, p<0.01).

Discussion

The current data provide evidence that symptoms of dissociation are extremely common in healthy humans

experiencing acute, highly intense stress. Further, such symptoms are significantly associated with the way in which a person has responded to, or has interpreted, a previously experienced traumatic event. The experience of “fearing for one’s life” significantly influenced the degree to which subjects experienced symptoms of dissociation before and during the stress of survival school. Dissociation was least prevalent in Special Forces soldiers—considered “stress hardy” by the U.S. Army—and was most prevalent in general infantry soldiers. Because Special Forces soldiers, as a group, tended to report more early trauma, yet dissociated the least under stress, they may represent individuals who have been “stress inoculated” by previous life events. Finally, symptoms of dissociation accounted for 41% of the variance in the physical health problems reported by subjects after stress exposure. This finding supports the idea that somatic distress and psychological adaptation to stress are linked.


The fact that acute stress-induced symptoms of dissociation were extremely common provides evidence that realistic military stress produces high levels of dissociation in psychologically healthy individuals. Indeed, the highly similar poststress dissociation scale scores in study 1 and study 2 indicate that the majority, and not the minority, of subjects experienced dissociation under stress. These findings suggest that the studies describing a “causal link” between peritraumatic dissociation and PTSD must be viewed with caution. Additional research is needed to clarify whether an etiological link between dissociation and PTSD exists.

General infantry soldiers experienced significantly more dissociative symptoms compared to Special Forces soldiers. This finding is consistent with the observations of other researchers that a large number of cases of combat-related stress reactions and subsequent PTSD arise in military personnel who are not front-line, elite, combat arms soldiers (25).

In study 2, individuals with the highest levels of dissociation were those who reported intense subjective responses to a previous trauma experience (“feared for their life”). These subjects reported higher levels of dissociation in the relatively mild classroom setting. However, when assessed after the high stress phase of the training, this group split into two subgroups: one that experienced an increase in dissociation and another that experienced low levels of dissociation. The subjects with lower levels of dissociation during stress were Special Forces soldiers. Although they had reported fearing for their life in response to a previous trauma, they exhibited the lowest increase of symptoms of dissociation in response to the actual stress of the experiential phase of survival school. By contrast, general infantry soldiers who reported fearing for their life in a previous trauma exhibited the greatest increase in dissociation in response to survival school stress.

One possible explanation is that exposure to early trauma or exposure to previous military training allowed

FIGURE 1. Scores on the Clinician-Administered Dissociative States Scale Before and After Exposure to Uncontrollable Stress^a in General Infantry Soldiers and Special Forces Soldiers With and Without Previous Exposure to Perceived Threat to Life



^a Uncontrollable stress presented by the experiential phase of survival training course, during which soldiers were subject to semi-starvation, sleep deprivation, lack of control over personal hygiene, and external control over movement, social contact, and communication.

^b Significant difference between soldiers with and without previous exposure to perceived threat to life ($t=3.2$, $df=38$, $p<0.003$).

^c Significant difference between soldiers with and without previous exposure to perceived threat to life ($t=3.6$, $df=38$, $p<0.001$).

the Special Forces soldiers to develop enhanced resistance to (or tolerance of) high-intensity stress. This phenomenon would be consistent with the concept of stress toughening (26). A second possible explanation is that individuals prone to higher levels of stress-induced dissociation may be screened out in the program used to select soldiers for the Special Forces.

Third, an individual’s formulation of his experience after a previous trauma may determine his subsequent responses to stress (27, 28). Individuals who form an appraisal of their experience that is nonalarming may be more stress tolerant. The relationship between postevent narratives, arousal, and responses to stress remains to be explored (29).

In both study 1 and study 2, Special Forces and general infantry soldiers differed significantly on poststress scores on Clinician-Administered Dissociative States Scale items that referred to an altered sense of one’s body, the environment, and other people. Previous investigation has shown that Special Forces subjects exhibited significantly higher plasma levels of neuropeptide Y in response to survival school stress (20). Neuropeptide Y has been shown to exhibit antianxiety and antistress properties, as well as an ability to enhance memory, alertness, and perception during stress (30, 31). It is possible that the differences in scores on the dissociation scale items are linked to an en-

hanced capacity for neuropeptide Y release during stress in Special Forces soldiers.

There was a significant and positive correlation between perceived health and dissociation. Dissociation accounted for 41% of the variance in somatic complaints. The presence of empirical, prospective evidence linking physical distress and dissociation is noteworthy and underscores the degree to which psychological adaptation to uncontrollable stress is linked to physical well-being (21, 32). These data highlight the need for medical professionals to assess concurrent psychological stress in patients being evaluated for physical complaints (33–35). Recent evidence has shown a significant, negative relationship between the severity of psychological symptoms of dissociation and the level of stress-induced release of neuropeptide Y. Although additional research is needed to clarify this issue, the current data raise the possibility that neuropeptide Y may also play a role in certain somatic complaints (20, 36).

This study has several limitations. First, due to the inherent restrictions of a naturalistic study, two separate groups of subjects participated in the study. In addition, subjects completed the Clinician-Administered Dissociative States Scale after being removed from the stressful situation, and not during the stressful situation. The degree to which the time elapsed since stress exposure may have influenced the subjective ratings is not known. For example, it is possible that the Special Forces and general infantry subjects experienced symptoms of dissociation to an equal degree during stress exposure, but that the Special Forces soldiers did not report these symptoms at the post-stress assessment. Such differences in reporting may be due to a personality style (or sense of machismo) that minimizes distress or discomfort—a style often seen in law enforcement personnel. It is unlikely that a reporting bias satisfactorily accounts for the current data since the Special Forces and the general infantry soldiers did not differ in their subjective reports of distress and anxiety during the training, as measured by Subjective Units of Distress Scales (20, 21). In addition, the Special Forces soldiers, as a group, tended to report early trauma on the Brief Trauma Questionnaire. Neither of these findings suggests that these subjects tended to underreport distressing symptoms. Further, our previous reports on the neuroendocrine responses of Special Forces soldiers showed that they had less overall stress-induced release of cortisol during this training than general infantry soldiers. Thus, the subjective and objective data both point to a diminished distress response in the Special Forces subjects compared to the general infantry subjects.

Another limitation is that this study included a combination of physical stressors (physical fatigue, starvation, sleep deprivation, etc.), environmental pressures (e.g., exposure to relative extremes of heat and cold), and psychological stress. Each of these elements may have contributed to participants' responses on the Clinician-

Administered Dissociative States Scale. Thus, it is not possible to determine which symptoms of dissociation were elicited by psychological stress alone. However, survival school stress is considered to represent a reasonable analog of the combination of physical and psychological stress experienced in combat situations. The use of an ethologically valid experience for soldiers increases the likelihood that the data can enhance our understanding of combat-related peritraumatic dissociation (37). Finally, the Health Problem List data were collected in a randomly selected group of participants. Although the significance of the regression analysis of the health problem data is noteworthy, given the small number of subjects, these findings may be influenced by the overall homogeneity of the military population and may not be representative of the civilian population. Nevertheless, military populations are routinely exposed to high-intensity stress, and these data underscore the degree to which psychological and physical responses may be linked.

In summary, symptoms of dissociation were common in healthy military subjects exposed to high-intensity stress. Individuals with a history of perceived threat to life in response to a traumatic event reported more symptoms of dissociation before experiencing the stress of survival training. Special Forces soldiers experienced significantly fewer symptoms of dissociation during stress, compared to general infantry soldiers. Symptoms of dissociation explained 41% of the variance in subjects' physical health complaints after stress exposure.

Received March 7, 2000; revision received Oct. 12, 2000; accepted Dec. 20, 2000. From the Department of Psychiatry, Yale University School of Medicine and Veterans Affairs Connecticut Healthcare System, West Haven, Conn.; JFK Special Warfare Training Center and School, Fort Bragg, N.C.; and the National Center for PTSD, White River Junction, Vt. Address reprint requests to Dr. Morgan, National Center for PTSD (116A), VA Connecticut Healthcare System, 950 Campbell Ave., West Haven, CT, 06516; charles.a.morgan@yale.edu (e-mail).

References

1. Kessler RC, Sonnega E, Bromet A, Bromet E, Hughes M, Nelson CB: Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995; 52:1048–1060
2. Breslau N, Davis GC, Andreski P, Peterson E: Traumatic events and posttraumatic stress disorder in an urban population of young adults. *Arch Gen Psychiatry* 1991; 48:216–222
3. Davidson JRT, Hughes D, Blazer D, Dan G, George LK: Posttraumatic stress disorder in the community: an epidemiological study. *Psychol Med* 1991; 21:1–9
4. Kulka RA, Schlenger WE, Fairbank JA, Hough RL, Jordan BK, Marmar CR, Weiss DS: Trauma and the Vietnam War Generation: Report of Findings From the National Vietnam Veterans Readjustment Study. New York, Brunner/Mazel, 1990
5. Breslau N, Chilcoat HD, Kessler RC, Davis GC: Previous exposure to trauma and PTSD effects of subsequent trauma: results from the Detroit Area Survey of Trauma. *Am J Psychiatry* 1999; 156:902–907
6. Duncan RD, Saunders BE, Kilpatrick DG, Hanson RF, Resnick HS: Childhood physical assault as a risk factor for PTSD, depres-

- sion, and substance abuse: findings from a national survey. *Am J Orthopsychiatry* 1996; 66:437-448
7. Freedy JR, Resnick HS, Kilpatrick DG, Dansky BS, Tidwell RP: The psychological adjustment of recent crime victims in the criminal justice system. *J Interpers Violence* 1994; 9:450-468
 8. Shalev AY, Peri T, Canetti L, Schreiber S: Predictors of PTSD in injured trauma survivors: a prospective study. *Am J Psychiatry* 1996; 153:219-225
 9. Spiegel D, Hunt T, Dondershine HE: Dissociation and hypnotizability in posttraumatic stress disorder. *Am J Psychiatry* 1988; 145:301-305
 10. Cardeña E, Spiegel D: Dissociative reactions to the San Francisco Bay Area earthquake of 1989. *Am J Psychiatry* 1993; 150:474-478
 11. Holen A: The North Sea oil rig disaster, in *International Handbook of Traumatic Stress Syndromes*. Edited by Wilson JP, Raphael B. New York, Plenum, 1993, pp 471-478
 12. Carlson EB, Rosser-Hogan R: Trauma experiences, posttraumatic stress, dissociation, and depression in Cambodian refugees. *Am J Psychiatry* 1991; 148:1548-1551
 13. Bremner JD, Southwick S, Brett E, Fontana A, Rosenheck R, Charney DS: Dissociation and posttraumatic stress disorder in Vietnam combat veterans. *Am J Psychiatry* 1992; 149:328-332
 14. Bremner JD, Brett E: Trauma-related dissociative states and long-term psychopathology in posttraumatic stress disorder. *J Trauma Stress* 1997; 10:37-49
 15. Koopman C, Classen C, Spiegel D: Predictors of posttraumatic stress symptoms among survivors of the Oakland/Berkeley, Calif, firestorm. *Am J Psychiatry* 1994; 151:888-894
 16. Marmar CR, Weiss DS, Schlenger WE, Fairbank JA, Jordan BK, Kulka RA, Hough RL: Peritraumatic dissociation and posttraumatic stress in male Vietnam theater veterans. *Am J Psychiatry* 1994; 151:902-907
 17. Marmar CR, Weiss DS, Metzler TJ, Delucchi KL, Best SR, Wentworth KA: Longitudinal course and predictors of continuing distress following critical incident exposure in emergency services personnel. *J Nerv Ment Dis* 1999; 187:15-22
 18. Farley M, Keaney JC: Physical symptoms, somatization, and dissociation in women survivors of childhood sexual assault. *Women Health* 1997; 25:33-45
 19. Schnurr PP, Spiro A III: Combat exposure, posttraumatic stress disorder symptoms, and health behaviors as predictors of self-reported physical health in older veterans. *J Nerv Ment Dis* 1999; 187:353-359
 20. Morgan CA III, Wang S, Southwick SM, Rasmusson A, Hauger R, Charney DS: Plasma neuropeptide-Y in humans exposed to acute uncontrollable stress. *Biol Psychiatry* 2000; 47:902-909
 21. Morgan CA III, Wang S, Mason J, Hazlett G, Fox P, Southwick SM, Charney DS, Greenfield G: Hormone profiles in humans experiencing military survival training. *Biol Psychiatry* 2000; 47:891-901
 22. Bremner JD, Krystal JH, Putnam FW, Southwick SM, Marmar C, Charney DS, Mazure CM: Measurement of dissociative states with the Clinician-Administered Dissociative States Scale (CADSS). *J Trauma Stress* 1998; 11:125-136
 23. Schnurr PP, Vieilhauer MJ, Weathers F, Findler M: *The Brief Trauma Questionnaire*. White River Junction, Vt, National Center for PTSD, 1999
 24. Schinka JA: *Health Problems Checklist*. Odessa, Fla, Psychological Assessment Resources, 1989
 25. Belenky G, Noy S, Solomon A: Battle stress, morale, cohesion, combat effectiveness, heroism, and psychiatric casualties: the Israeli experience, in *Contemporary Studies in Combat Psychiatry: Contributions in Military Studies* 62. Edited by Belenky G. Westport, Conn, Greenwood Press, 1987, pp 11-20
 26. Dienstbier RA: Arousal and physiological toughness: implications for mental and physical health. *Psychol Rev* 1989; 96:84-100
 27. Meichenbaum D, Fitzpatrick D: A constructivist narrative perspective on stress and coping: stress inoculation and applications, in *Handbook of Stress: Theoretical and Clinical Aspects*, 2nd ed. Edited by Goldberger L, Breznitz S. New York, Free Press, 1993, pp 706-723
 28. Meichenbaum D, Fong GT: How individuals control their own minds: a constructive narrative perspective, in *Handbook of Mental Control*. Edited by Wegner DM, Pennebaker JW. Englewood Cliffs, NJ, Prentice-Hall, 1993, pp 473-490
 29. Chrousos GP, Gold PW: The concepts of stress and stress system disorders: overview of physical and behavioral homeostasis. *JAMA* 1992; 267:1244-1252
 30. Broqua P, Wettstein JG, Rocher MN, Gauthier-Martin B, Junien JL: Behavioral effects of neuropeptide Y receptor agonists in the elevated plus-maze and fear-potentiated startle procedures. *Behav Pharmacol* 1995; 6:215-222
 31. Heilig M, Koob GF, Britton KT: Anxiolytic-like effect of neuropeptide Y (NPY), but not other peptides, in an operant conflict test. *Regul Pept* 1992; 41:65-69
 32. Herschbach P, Henrich G, von Rad M: Psychological factors in functional gastrointestinal disorders: characteristics of the disorder or of the illness behavior? *Psychosom Med* 1999; 61:148-153
 33. Post D, Miller K: Use of the health history as a psychiatric screening tool. *J Am Board Fam Pract* 1998; 11:452-458
 34. Russo J, Katon W, Sullivan M, Clark M, Buckwald D: Severity of somatization and its relationship to psychiatric disorders and personality. *Psychosomatics* 1994; 35:546-556
 35. Kroenke K, Spitzer RL, Williams JB, Linzer M, Hahn SR, deGruy FV III, Brody D: Physical symptoms in primary care: predictors of psychiatric disorders and functional impairment. *Arch Fam Med* 1994; 3:774-779
 36. Pruessner JG, Hellhammer DH, Kirschbaum C: Burnout, perceived stress, and cortisol responses to awakening. *Psychosom Med* 1999; 61:197-204
 37. Nijenhuis ERS, Vanderlinden J, Spinhoven P: Animal defensive reactions as a model for trauma-induced dissociative reactions. *J Trauma Stress* 1998; 11:243-260