

# Symptoms of Posttraumatic Stress Disorder and Borderline Personality Disorder in Veterans of Operation Desert Storm

Seth R. Axelrod, Ph.D.

C. Andrew Morgan III, M.D.

Steven M. Southwick, M.D.

**Objective:** The present report is part of a follow-along investigation focusing on the evolution of trauma-related symptoms in veterans of Operation Desert Storm. The goal of the current report was to examine three hypotheses on the relationship between severity of war-related trauma, symptoms of posttraumatic stress disorder (PTSD), and symptoms of borderline personality disorder with a mixed retrospective/prospective design.

**Method:** Ninety-four National Guard reservists completed self-administered measures of combat-related trauma, PTSD symptoms, and borderline personality disorder features after their Gulf War duty.

**Results:** Consistent with study hypotheses, prewar features of borderline per-

sonality disorder predicted variability in postwar PTSD symptoms beyond that predicted by combat exposure, combat exposure predicted variability in postwar features of borderline personality disorder, and PTSD severity assessed shortly after combat exposure accounted for additional variability in subsequent features of borderline personality disorder.

**Conclusions:** Taken together, the present findings suggest that trauma, symptoms of PTSD, and features of borderline personality disorder are related to one another in a complex fashion that may exceed simple linear models. Clinical and research implications for the relationships among trauma, PTSD, and borderline personality disorder are discussed.

(*Am J Psychiatry* 2005; 162:270–275)

A number of studies have reported high rates of borderline personality disorder or features in combat veterans with posttraumatic stress disorder (PTSD) (1–4). There are several possible explanations for these high rates. First, preexisting features of borderline personality disorder may serve as a risk factor for the development of PTSD. Gunderson and Sabo (5) suggest that individuals with borderline personality disorder have limited resources for dealing with traumatic events, which leaves them more vulnerable toward developing PTSD. Second, it may be that trauma, in and of itself, causes enduring changes in character. This explanation is consistent with a large body of evidence showing high rates of childhood trauma in adults with borderline personality disorder (6–9). In fact, Herman and van der Kolk (10) view borderline personality disorder, like PTSD, as a stress-related disorder. Third, it could be that borderline features develop in response to living with the core symptoms of PTSD rather than to trauma per se (1). For example, nightmares and sleep deprivation can lead to irritability, intense anger, and affective instability—symptoms commonly associated with borderline personality disorder.

Past efforts to clarify the nature or direction of relationships among trauma, PTSD, and borderline personality disorder have been hindered by limitations in research methods. To date, cross-sectional approaches have been

used to investigate the comorbidity of PTSD and borderline personality disorder, and as a result, issues regarding causality have not been adequately addressed. Further, investigation of traumatic stressors in the etiology of borderline personality disorder has primarily relied on retrospective accounts from adult clinical samples (6–9), which may be affected by subject selection and potential reporting biases (11, 12). More recent investigations have used prospective designs for examining trauma in the development of borderline personality disorder (13, 14) but have not included assessment of pretrauma personality functioning for comparison.

The present report is part of a follow-along investigation focusing on the evolution of trauma-related symptoms in veterans of Operation Desert Storm. Desert Storm veterans recruited from Connecticut National Guard units filled out self-administered questionnaires at multiple time points after their return from the Gulf. The goal of the current report was to examine the relationship among the severity of war-related trauma, symptoms of PTSD, and borderline personality disorder features in veterans of Desert Storm.

This report uses a mixed retrospective/prospective design to address some of the limitations of prior research. Pretrauma borderline personality disorder features were assessed retrospectively and used to predict postwar PTSD

symptoms, and PTSD symptoms measured shortly after return from the Gulf were used to predict subsequent borderline personality disorder features. The study hypotheses were as follows:

Hypothesis 1: prewar borderline personality disorder features will account for variability in postwar PTSD symptoms (beyond the effects of combat exposure) (5).

Hypothesis 2: 1) combat exposure will predict increased postwar borderline personality disorder features (beyond variability that can be accounted for by prewar borderline personality disorder features), and 2) younger age at combat exposure will predict additional variability in postwar borderline personality disorder features (15–17).

Hypothesis 3: combat-related PTSD symptoms will predict postwar borderline personality disorder features (beyond variability accounted for by hypothesis 2) (1).

## Method

### Study Participants

Ninety-four participants were part of a prospective study designed to examine the longitudinal course of trauma-related symptoms in veterans of the Gulf War. The participants were 73.4% male, their ages ranged from 19 to 53 years (mean=29.0, SD=9.5), and they attended an average of 13.7 years of school (SD=1.8). Their racial/ethnic background was 78.7% Caucasian, 10.6% African American, 6.4% Hispanic American, and 4.3% Asian American. After a complete description of the study to the participants, written informed consent was obtained.

### Procedures

Much of the current methodology has been used in a number of previous reports involving Desert Storm veterans (18, 19). The participants completed a battery of self-administered questionnaires at 1 month and at 6 months after their return from the Gulf. The present study focuses on three of these questionnaires, including the Desert Storm Trauma Questionnaire (18), that were completed 1 month after their return from the Gulf, a PTSD symptom scale (18) completed at 1 month and again at 6 months after their return, and the Personality Diagnostic Questionnaire (20), which was completed 6 months after their return. For the purposes of this study, we instructed participants to apply the standard Personality Diagnostic Questionnaire instructions to each question for two time periods: the 6 months before their deployment to the Gulf and the 6 months after their return from the Gulf. The participants recorded these ratings on the same page as the Personality Diagnostic Questionnaire items by entering their responses in columns labeled “6 Months Before the War” and “Past 6 Months.”

### Measures

The Desert Storm Trauma Questionnaire (18) is composed of 19 items dealing with stressors that were experienced by many Desert Storm personnel. These include extreme threats to personal safety, seeing others killed or wounded, the death of a close friend, sitting with the dying, being stationed close to enemy lines, and witnessing bizarre disfigurement of bodies as a result of wounds.

The total Desert Storm Trauma Questionnaire score is the number of positive responses. The PTSD symptom scale (18) is based on the DSM-III-R criteria (largely unchanged in DSM-IV). The degree of severity of each symptom during the preceding month was rated on a scale in which 0=none, 1=slight, 2=moderate, 3=considerable, and 4=extreme. A total PTSD score was calculated by summing the ratings for each symptom.

The Personality Diagnostic Questionnaire (20) was designed to assess the 11 personality disorders included in the DSM-III. Psychometric data for the Personality Diagnostic Questionnaire Revised for DSM-III-R suggest that the personality disorder scales have good 3-month test-retest reliability (21) and excellent sensitivity with respect to a semistructured interview but only modest specificity (22). For these reasons, the Personality Diagnostic Questionnaire is best used for screening personality disorders or for studying personality disorders dimensionally. At the suggestion of the Personality Diagnostic Questionnaire's author (Hyler, personal communication), we measured DSM-IV borderline personality disorder features by selecting Personality Diagnostic Questionnaire items that best captured DSM-IV borderline personality disorder (reflecting the changes made in the borderline personality disorder criteria since DSM-III). We identified Personality Diagnostic Questionnaire items to represent eight of the nine DSM-IV borderline personality disorder criteria (all but frantic efforts to avoid abandonment). In order to avoid inflated correlation of borderline personality disorder and PTSD because of similarity in diagnostic criteria, we excluded Personality Diagnostic Questionnaire content related to anger. Borderline personality disorder features were then measured as the sum of the seven criteria assessed (scores ranging from 0 to 7).

### Data Analysis

The three study hypotheses were examined with hierarchical regression modeling. Hypothesis 1 was examined by including PTSD symptoms as the dependent variable and entering the Desert Storm Trauma Questionnaire score and borderline features in separate steps as independent variables. Support for hypothesis 1 would be indicated if borderline features significantly contributed to PTSD symptom variability beyond that accounted for by the Desert Storm Trauma Questionnaire score. This analysis was conducted for PTSD symptoms at 1 month and at 6 months.

Hypothesis 2 was examined by treating postwar borderline personality disorder features as the dependent variable. Here prewar borderline personality disorder features were entered as an independent variable in the first step, the Desert Storm Trauma Questionnaire score was entered as a second step, and age at combat exposure was entered as a third step. Hypothesis 2 would be supported if Desert Storm Trauma Questionnaire score variability accounted for postwar borderline personality disorder features beyond variability that could be accounted for by prewar borderline personality disorder features. Additional support would be provided if a younger age (at the time of the trauma) was associated with borderline personality disorder features.

Hypothesis 3 is an extension of Hypothesis 2 and was examined by entering PTSD symptoms at 1 month as an additional independent variable in a fourth step. Hypothesis 3 would be supported to the extent that PTSD symptoms at 1 month accounted for variability in borderline personality disorder features at 6 months postwar beyond the variability that could be accounted for by prewar borderline personality disorder features, Desert Storm Trauma Questionnaire score, and age.

## Results

Desert Storm Trauma Questionnaire scores ranged from 0 to 11, with a mean of 3.3 (SD=2.8). The PTSD symptom scale had a mean of 8.7 (SD=9.7) 1 month after the veterans' return from the Gulf and 12.0 (SD=12.1) at 6 months after the return. Borderline personality disorder features for the 6 months before their return from the Gulf ranged from 0 to 5, with a mean of 1.5 (SD=1.3), and borderline personality disorder features for the 6 months after their

**TABLE 1. Regression Models Predicting 1-Month and 6-Month Symptoms of Posttraumatic Stress Disorder in Veterans of Operation Desert Storm**

R <sup>2</sup>	ΔR <sup>2</sup>	F	df	p
1-month symptoms (N=94)				
Step 1: Desert Storm Trauma Questionnaire score	0.24	29.51	1, 92	<0.001
Step 2: prewar borderline features	0.01	1.77	1, 91	0.19
6-month symptoms (N=90)				
Step 1: Desert Storm Trauma Questionnaire score	0.30	38.31	1, 88	<0.001
Step 2: prewar borderline features	0.05	6.74	1, 87	<0.02

return to the Gulf ranged from 0 to 6, with a mean of 1.7 (SD=1.5).

Table 1 presents the regression models evaluating hypothesis 1. Desert Storm Trauma Questionnaire score accounted for 24% of the 1-month PTSD symptom variability ( $p<0.001$ ) and 30% of the 6-month variability ( $p<0.001$ ). Prewar borderline personality disorder features accounted for an additional 1% of the 1-month PTSD symptom variability ( $p=0.19$ ) and 5% of the 6-month variability ( $p<0.02$ ), beyond the variability accounted for by the Desert Storm Trauma Questionnaire.

Table 2 presents the regression models examining hypotheses 2 and 3. Prewar borderline personality disorder features were entered in the first step of the model and accounted for 40% of the variability of the postwar borderline personality disorder features ( $p<0.001$ ). To analyze hypothesis 2, Desert Storm Trauma Questionnaire score was entered in a second step and accounted for 10% of the variability in postwar borderline personality disorder features ( $p<0.001$ ), and age was entered in a third step and accounted for 3% of the variability ( $p<0.02$ ). Finally, to analyze hypothesis 3, 1-month PTSD symptoms were entered in a fourth step, accounting for an additional 11% of variability ( $p<0.001$ ).

## Discussion

In the present report on Operation Desert Storm veterans, prewar features of borderline personality disorder predicted variability in postwar PTSD symptoms beyond that predicted by combat exposure, combat exposure predicted variability in postwar features of borderline personality disorder, and PTSD severity assessed shortly after combat exposure accounted for additional variability in subsequent features of borderline personality disorder. The findings support each of the three study hypotheses and point to a complex relationship among trauma, PTSD symptoms, and borderline personality disorder features.

Prewar borderline personality disorder features accounted for a significant amount of variance in PTSD symptoms at 6 months postwar (5%) but did not significantly account for PTSD symptoms at 1 month postwar. The finding with regard to 6-month PTSD is consistent with Gunderson and Sabo's suggestion (5) that individuals with borderline personality lack adequate coping mecha-

**TABLE 2. Regression Model Predicting Postwar Borderline Personality Disorder in 94 Veterans of Operation Desert Storm**

R <sup>2</sup>	ΔR <sup>2</sup>	F	df	p
Step 1: prewar borderline features	0.40	60.50	1, 92	<0.001
Step 2: Desert Storm Trauma Questionnaire score	0.10	17.35	1, 91	<0.001
Step 3: age	0.03	6.44	1, 90	<0.02
Step 4: 1-month symptoms of posttraumatic stress disorder	0.11	27.09	1, 89	<0.001

nisms and are therefore vulnerable to posttraumatic stress reactions after trauma. It is also possible that individuals with borderline personality disorder features, such as dangerous impulsivity and uncontrolled anger, are more likely to engage in situations where they are exposed to trauma, putting them at increased risk for developing PTSD. However, our data did not find a significant correlation between prewar borderline personality disorder features and combat exposure ( $r=0.12$ ,  $p=0.26$ ). Other personality features that have been reported as risk factors for the development of PTSD include neuroticism, extraversion, external locus of control, and poor self-esteem (23, 24). Neuroticism, external locus of control, and poor self-esteem share variability with borderline personality disorder features, but extraversion does not (25). Additional research may identify other personality disorder traits that create a vulnerability to PTSD.

The second hypothesis was supported in that level of Desert Storm trauma exposure accounted for 10% of the variability in postwar borderline personality disorder features beyond the variability accounted for by prewar borderline personality disorder features. Exposure to early life trauma, such as childhood physical, sexual, and emotional abuse, has long been considered a risk factor for the development of borderline personality and has been supported by prospective research (13, 14). However, traumatic exposure during late adolescence and early adulthood has not typically been associated with symptoms of this disorder.

The third hypothesis was also supported because PTSD symptoms at 1 month accounted for 11% of the variability in borderline personality disorder features at 6 months above and beyond the contributions from prewar borderline personality disorder features, war trauma (Desert Storm Trauma Questionnaire score), and age. Herman (15) has suggested that the DSM-IV "simple" PTSD symptoms of intrusiveness, avoidance, and hyperarousal are typical stress responses to acute traumatic experiences but that when individuals experience prolonged or repeated trauma, they also develop "complex PTSD" responses (also called disorders of extreme stress not otherwise specified), including borderline personality disorder-like symptoms. She described the nature of prolonged stress, such as being subjugated to coercion, and the adjustments that individuals make that result in complex PTSD symptoms. Our data indicate that the development of "simple"

PTSD symptoms in and of themselves may create challenges that require personality adjustments. "Simple" PTSD symptoms, such as hypervigilance to cues, emotional detachment, and chronically disturbed sleep, are often highly disruptive and debilitating. Living with these symptoms may lead to a number of experiences commonly associated with borderline personality disorder, including marked affective instability, unstable and intense interpersonal relationships, and inappropriate intense anger. We believe that living with "simple" PTSD symptoms provides an additional component to the etiology of complex PTSD, as described by Herman (15).

The current findings suggest that adulthood traumatic experiences and posttraumatic stress sequelae may contribute to the development of borderline personality disorder features. The onset of borderline personality disorder symptoms in adulthood is consistent with two diagnostic categories that were considered for the DSM-IV but not included. In addition to complex PTSD/disorders of extreme stress not otherwise specified, which was ultimately added to the associated features text of DSM-IV PTSD (26, 27), the ICD-10 diagnosis of enduring personality change after catastrophic experience was considered for DSM-IV inclusion (28). If the current findings are extended by additional research linking adult trauma exposure to borderline personality disorder-like personality changes that cause clinically significant distress or impairment, then further consideration should be given to adding these trauma-related diagnostic categories to DSM-V.

Common neurobiological features related to affect dysregulation, impulsivity, and stress reactivity may contribute to the high rate of co-occurrence of borderline personality disorder and PTSD. Examples of this shared pathophysiology include decreased platelet  $\alpha_2$ -adrenergic receptor number (29, 30), hypothalamic-pituitary-adrenal axis hyperresponsivity to pharmacologic challenge (31, 32), elevated CSF corticotrophin-releasing hormone (33, 34), reduced hippocampal volume (35, 36), and exaggerated amygdala activation in response to stressful stimuli (37, 38). Of course, further research will be necessary to identify potential relationships among neurobiological alterations, symptom development, and the possible direction of influence between borderline personality disorder and PTSD.

There are a number of limitations in the current study. First, participants were non-treatment-seeking and did not exhibit high rates of psychopathology (18). Therefore, the relationships among trauma, PTSD symptoms, and borderline personality disorder features may not replicate with treatment-seeking patients. On the other hand, given the low rates of pathology, it is unlikely that the self-reports of these participants would have been compromised by disability or compensation claims, and in fact, we were unaware of any participants pursuing such claims, and no participants requested to have study assessments released for such purposes. Also, given that the study hypotheses were supported within the restricted range of pathology of

this sample, it would be reasonable to expect the relationships to be maintained in samples with greater variability. Future prospective investigations with larger military samples are necessary to examine these relationships in participants that develop full PTSD and/or borderline personality disorder conditions.

Second, the current data are limited to self-report measures of combat trauma, PTSD symptoms, and borderline personality disorder features. Objective measurement of combat experience is not available to assess the criterion validity of the Desert Storm Trauma Questionnaire self-report, and there is some indication that self-reported accounts of combat exposure are not perfectly reliable. In a previous study that used many of the same participants from the current study (39), we found that self-reported accounts of combat exposure did not remain perfectly consistent across time, particularly when attending to responses on individual exposure items. However, the participants' total number of types of exposure (or Desert Storm Trauma Questionnaire total scores) was fairly consistent, changing on average by less than one type of exposure (39). The present analyses relied upon Desert Storm Trauma Questionnaire total scores and are therefore not likely to have been significantly affected by inconsistent reporting of combat exposure.

Third, the Personality Diagnostic Questionnaire scales have demonstrated low diagnostic thresholds, thus overdiagnosing personality disorders (22). Therefore, it would not be appropriate to use the Personality Diagnostic Questionnaire to estimate rates of borderline personality disorder in this group. However, low diagnostic thresholds are advantageous for examining dimensional relationships of personality disorder features in healthy samples because more stringent criteria would not yield sufficient variability for use in a nonclinical sample.

Fourth, prewar and postwar borderline personality disorder features, as measured by Personality Diagnostic Questionnaire items, correlated at 0.62. Had these measures been given prospectively, a correlation of this magnitude might have suggested only modest test-retest reliability. However, each Personality Diagnostic Questionnaire item was completed by the participants for the two intervals at the same time (literally marked on the same page), indicating that any difference in scores was a deliberate reflection of the participants' change in self-perception between the two intervals and not due to a lack of reliability in the self-report measure. On the other hand, the retrospective account of prewar personality functioning is a significant limitation of this data. It will be important to conduct prospective assessments of prewar and postwar personality functioning in future investigations, although such investigations will obviously require significant resources.

Finally, given the nature of the military strategies employed in Operation Desert Storm (i.e., strong reliance on missiles and air assaults and limited ground confronta-



tion), it might be presumed that the degree of combat exposure would have been insufficient to result in posttraumatic stress reactions. Nonetheless, most participants were confronted with multiple combat-related stressors. Participants in this study were frequently exposed to incoming SCUD missile attacks, and some participants were reportedly stationed as guards at the entrance to their compound and were prevented from taking refuge in bunkers during missile attacks. In addition, some soldiers were exposed to enemy small-arms fire (many participants from one unit witnessed the deaths of a physician and a nurse), and several participants reported seeing burned and charred bodies and flattened corpses.

The findings of this study have a number of potential implications. First, for individuals choosing or hiring for professions that involve a high risk of becoming traumatized, such as military or law enforcement, it is important to consider that features of personality may serve as potential risk factors for developing PTSD. Second, when diagnosing borderline personality disorder features in adulthood, clinicians should consider the contribution of adulthood trauma as well as childhood trauma. Further, when one is treating traumatized individuals with PTSD and comorbid features of borderline personality disorder, lessening the symptoms of PTSD may, over time, help to reduce coexisting borderline personality disorder features. Alternatively, targeting borderline personality disorder-related problems, such as affective dysregulation and impulsivity (e.g., reference 40), might be necessary before patients will be able to fully benefit from treatments focused on resolving PTSD symptoms (e.g., reference 41). Finally, the presence of personality disorder features can alter the way in which PTSD patients are viewed and/or treated by clinicians, researchers, and disability claim adjusters. For example, patients with symptoms of borderline personality disorder may be avoided or labeled as problematic, and traumatized individuals may be denied disability claims, despite the fact that trauma appears to contribute to borderline personality disorder features, as well as PTSD.

Taken together, the present findings suggest that trauma, symptoms of PTSD, and features of borderline personality disorder are related to one another in a complex fashion that may exceed simple linear models. The features of borderline personality disorder appear to be both a risk factor for the development of PTSD symptoms and a potential consequence of living with PTSD symptoms. In addition, it appears that trauma can contribute to changes in personality, even in adulthood. Future prospective investigations are necessary to clarify the role of military trauma and other adult traumas in the development of borderline personality disorder symptoms in adulthood. Longitudinal investigations may reveal a positive feedback loop between PTSD symptoms and borderline personality disorder features that helps to maintain both conditions over time.

Presented in part at the 155th annual meeting of the American Psychiatric Association, Philadelphia, May 18–23, 2002; and at the 110th annual meeting of the American Psychological Association, Chicago, Aug. 22–25, 2002. Received Feb. 12, 2003; revision received Jan. 30, 2004; accepted April 23, 2004. From the Department of Psychiatry, Yale University School of Medicine; and the National Center for PTSD, VA Connecticut Healthcare System, Newington, Conn. Address correspondence and reprint requests to Dr. Axelrod, Yale University School of Medicine, 425 George St., New Haven, CT 06511; seth.axelrod@yale.edu (e-mail).

Supported by the National Center for PTSD and the Yale Borderline Personality Disorder Research Center.

The authors thank Paula Schnurr, Ph.D., and Martha Dillon, M.A., for their contributions.

## References

- Southwick SM, Yehuda R, Giller EL Jr: Personality disorders in treatment-seeking combat veterans with posttraumatic stress disorder. *Am J Psychiatry* 1993; 150:1020–1023
- Sherwood RJ, Funari DJ, Piekarski AM: Adapted character styles of Vietnam veterans with posttraumatic stress disorder. *Psychol Rep* 1990; 66:623–631
- Hyer L, Woods MG, Boudewyns PA: A three tier evaluation of PTSD among Vietnam combat veterans. *J Trauma Stress* 1991; 4:165–194
- Shea TM, Zlotnick C, Weisberg RB: Commonality and specificity of personality disorder profiles in subjects with trauma histories. *J Personal Disord* 1999; 13:199–210
- Gunderson JG, Sabo AN: The phenomenological and conceptual interface between borderline personality disorder and PTSD. *Am J Psychiatry* 1993; 150:19–27
- Herman J, Perry JC, van der Kolk BA: Childhood trauma in borderline personality disorder. *Am J Psychiatry* 1989; 146:490–495
- Ogata S, Silk KR, Goodrich S, Lohr NE, Westen D, Hill EM: Childhood sexual and physical abuse in adult patients with borderline personality disorder. *Am J Psychiatry* 1990; 147:1008–1013
- Paris J, Zweig-Frank H, Guzder J: Risk factors for borderline personality disorder in male outpatients. *J Nerv Ment Dis* 1994; 182:375–380
- Zanarini MC, Gunderson JG, Marino MG, Schwartz EO, Frankenburg FR: Childhood experiences of borderline patients. *Compr Psychiatry* 1989; 30:18–25
- Herman JL, van der Kolk BA: Traumatic antecedents of borderline personality disorder, in *Psychological Trauma*. Edited by van der Kolk BA. Washington, DC, American Psychiatric Press, 1987, pp 111–126
- Paris J: Childhood trauma as an etiological factor in the personality disorders. *J Personal Disord* 1997; 11:34–49
- Paris J, Zweig-Frank H: A critical review of the role of childhood sexual abuse in the etiology of borderline personality disorder. *Can J Psychiatry* 1992; 37:125–128
- Johnson JJ, Smailes EM, Cohen P, Brown J, Bernstein DP: Associations between four types of childhood neglect and personality disorder symptoms during adolescence and early adulthood: findings of a community-based longitudinal study. *J Personal Disord* 2000; 14:171–187
- Johnson JG, Cohen P, Brown J, Smailes EM, Bernstein DP: Childhood maltreatment increases risk for personality disorders during early adulthood. *Arch Gen Psychiatry* 1999; 56:600–606
- Herman JL: Complex PTSD: a syndrome in survivors of prolonged and repeated trauma. *J Trauma Stress* 1992; 5:377–391
- van der Kolk BA: The complexity of adaptation to trauma self-regulation, stimulus discrimination, and characterological development, in *Traumatic Stress*. Edited by van der Kolk BA, Al-

- exander C, McFarlane LW. New York, Guilford, 1996, pp 378–397
17. van der Kolk BA, Hostetler A, Herron N, Fisler R: Trauma and the development of borderline personality disorder. *Psychiatr Clin North Am* 1994; 17:715–730
18. Southwick SM, Morgan CA, Nagy LM, Bremner D, Nicolaou AL, Johnson DR, Rosenheck R, Charney DS: Trauma-related symptoms in veterans of Operation Desert Storm: a preliminary report. *Am J Psychiatry* 1993; 150:1524–1528
19. Southwick SM, Morgan CA III, Darnell A, Bremner D, Nicolaou AL, Nagy LM, Charney DS: Trauma-related symptoms in veterans of Operation Desert Storm: a 2-year follow-up. *Am J Psychiatry* 1995; 152:1150–1155
20. Hyler SE, Rieder RO, Williams JBW, Spitzer RL, Hendler J, Lyons M: The Personality Diagnostic Questionnaire: development and preliminary results. *J Personal Disord* 1988; 2:229–237
21. Trull TJ: Temporal stability and validity of two personality inventories. *Psychol Assess* 1993; 5:11–18
22. Hyler SE, Skodol AE, Kellman HD, Oldham JM, Rosnick L: Validity of the Personality Diagnostic Questionnaire, Revised: comparison with two structured interviews. *Am J Psychiatry* 1990; 147:1043–1048
23. Bremner JD, Southwick SM, Charney DS: Etiological factors in the development of posttraumatic stress disorder, in *Does Stress Cause Psychiatric Illness?* Edited by Mazure CM. Washington DC, American Psychiatric Press, 1995, pp 149–185
24. Schnurr PP, Friedman MJ, Rosenberg SD: Premilitary MMPI scores as predictors of combat-related PTSD symptoms. *Am J Psychiatry* 1993; 150:479–483
25. Watson DC: The relationship of self-esteem, locus of control, and dimensional models to personality disorders. *J Soc Behav Pers* 1998; 13:399–420
26. Roth S, Newman E, Pelcovitz D, van der Kolk B, Mandel F: Complex PTSD in victims exposed to sexual and physical abuse: results from the DSM-IV field trials. *J Trauma Stress* 1997; 10:539–555
27. van der Kolk BA, Roth S, Pelcovitz D, Mandel F: Complex PTSD: Results of the PTSD Field Trials for DSM-IV. Washington, DC, American Psychiatric Press, 1993
28. Shea MT: Enduring personality change after catastrophic experience, in *DSM-IV Sourcebook, Vol 2*. Edited by Widiger TA. Washington, DC, American Psychiatric Press, 1996, pp 849–860
29. Southwick SM, Yehuda R, Giller EL, Perry BD: Altered platelet alpha-2 adrenergic receptor binding sites in borderline personality disorder. *Am J Psychiatry* 1990; 147:1014–1017
30. Perry BD, Giller EL, Southwick SM: Altered platelet alpha 2-adrenergic binding sites in posttraumatic stress disorder. *Am J Psychiatry* 1987; 144:1511–1512
31. Yehuda R, Southwick SM, Krystal JH, Bremner D, Charney DS, Mason JW: Enhanced suppression of cortisol binding following dexamethasone administration in post-traumatic stress disorder. *Am J Psychiatry* 1993; 150:83–86
32. Rinne T, De Kloet ER, Wouters L, Goekoop J, De Rijk R, Van den Brink W: Fluvoxamine reduces responsiveness of HPA axis in adult female borderline personality disorder patients with a history of sustained childhood abuse. *Neuropsychopharmacology* 2003; 28:126–132
33. Lee RL, Coccaro EF: Childhood trauma and alterations in adult CRF in personality disorders, in 2002 Annual Meeting New Research Program and Abstracts. Washington, DC, American Psychiatric Association, 2002, number 468
34. Bremner JD, Licinio J, Darnell A, Krystal JH, Owens MJ, Southwick SM, Nemeroff CB, Charney DS: Elevated CSF corticotropin-releasing factor concentrations in posttraumatic stress disorder. *Am J Psychiatry* 1997; 154:624–629
35. Bremner JD, Randall P, Scott TM, Bronen RA, Seibyl JP, Southwick SM, Delaney RC, McCarthy G, Charney DS, Innis RB: MRI-based measurement of hippocampal volume in patients with combat-related posttraumatic stress disorder. *Am J Psychiatry* 1995; 152:973–981
36. Driessen M, Herrmann J, Stahl K, Zwaan M, Meier S, Hill A, Osterheider M, Petersen D: Magnetic resonance imaging volumes of the hippocampus and the amygdala in women with borderline personality disorder and early traumatization. *Arch Gen Psychiatry* 2000; 57:1115–1122
37. Rauch SL, Whalen PJ, Shin LM, McNerney SC, Macklin ML, Lasko NB, Orr SP, Pitman RK: Exaggerated amygdala response to masked facial stimuli in posttraumatic stress disorder: a functional MRI study. *Biol Psychiatry* 2000; 47:769–776
38. Herpertz SC, Dietrich TM, Wenning B, Krings T, Erberich SG, Willmes K, Thron A, Sass H: Evidence of abnormal amygdala functioning in borderline personality disorder: a functional MRI study. *Biol Psychiatry* 2001; 15:292–298
39. Southwick SM, Morgan CA III, Nicolaou AL, Charney DS: Consistency of memory for combat-related traumatic events in veterans of Operation Desert Storm. *Am J Psychiatry* 1997; 154:173–177
40. Linehan MM: *Cognitive Behavioral Treatment of Borderline Personality Disorder*. New York, Guilford, 1993
41. Foa EB, Keane TM, Friedman MJ (eds): *Effective Treatments for PTSD: Practice Guidelines From the International Society for Traumatic Stress Studies*. New York, Guilford, 2000