

8-Year Follow-Up of Patients Treated for Borderline Personality Disorder: Mentalization-Based Treatment Versus Treatment as Usual

Anthony Bateman, F.R.C.Psych.

Peter Fonagy, Ph.D., F.B.A.

Objective: This study evaluated the effect of mentalization-based treatment by partial hospitalization compared to treatment as usual for borderline personality disorder 8 years after entry into a randomized, controlled trial and 5 years after all mentalization-based treatment was complete.

Method: Interviewing was by research psychologists blind to original group allocation and structured review of medical notes of 41 patients from the original trial. Multivariate analysis of variance, chi-square, univariate analysis of variance, and nonparametric Mann-Whitney statistics were used to contrast the two groups depending on the distribution of the data.

Results: Five years after discharge from mentalization-based treatment, the men-

talization-based treatment by partial hospitalization group continued to show clinical and statistical superiority to treatment as usual on suicidality (23% versus 74%), diagnostic status (13% versus 87%), service use (2 years versus 3.5 years of psychiatric outpatient treatment), use of medication (0.02 versus 1.90 years taking three or more medications), global function above 60 (45% versus 10%), and vocational status (employed or in education 3.2 years versus 1.2 years).

Conclusions: Patients with 18 months of mentalization-based treatment by partial hospitalization followed by 18 months of maintenance mentalizing group therapy remain better than those receiving treatment as usual, but their general social function remains impaired.

(*Am J Psychiatry* 2008; 165:631–638)

The natural course of borderline personality disorder and its long-term outcome following treatment are uncertain (1). A number of well-characterized treatments for borderline personality disorder have been found in randomized, controlled trials to reduce suicidal acts, self-harm, impulsive behaviors, general psychopathology, and service use while improving affective control (2–7). More limited evidence exists from these trials for changes in depression, loneliness/emptiness, anger, and social and interpersonal function with little confirmation of sustained improvement in any of these domains. Follow-up after treatment was either absent or too short to assess final outcomes.

Naturalistic follow-along investigations report symptomatic improvement, particularly of impulsive symptoms, over a relatively short period of time but suggest that deficits in interpersonal and social function and vocational achievement (8) remain over the longer term (9, 10). But it is difficult to draw firm conclusions about either the natural or treated course of the disorder in the absence of an experimental design with well-defined interventions.

In the short term, controlled studies have found limited between-groups differences at 2 years after entrance into

treatment (6, 11, 12), implying that some treatments may achieve a more rapid natural remission. Longer-term follow-up studies suggesting that posttreatment differences are maintained have lacked adequate comparison groups (13, 14).

We reported 18-month (end of intensive treatment) and 36-month outcomes of patients treated for borderline personality disorder after random assignment to mentalization-based treatment by partial hospitalization or treatment as usual (15, 16). Mentalization-based treatment by partial hospitalization and treatment as usual for 18 months were well-characterized. Subsequent treatment was monitored. However, the mentalization-based treatment by partial hospitalization group continued to receive some outpatient group mentalizing treatment between 18 and 36 months. No treatment as usual patients received the experimental treatment during this 36-month period. Differences between groups found at the end of intensive treatment not only were maintained during 18–36 months but increased substantially. We attributed this to the rehabilitative processes stimulated by the initial mentalization-based treatment by partial hospitalization. But equally it might have been a result of the maintenance

This article is discussed in an editorial by Dr. Levy (p. 556).

outpatient group mentalizing treatment even though this group had considerably less treatment than the control group.

All mentalization-based treatment ended 36 months after entry into the study. We wanted to determine whether treatment gains were maintained over the subsequent 5 years, i.e., 8 years after random assignment. The primary outcome measure for this long-term follow-up study was the number of suicide attempts. But in light of the limited improvement related to social adjustment in follow-along studies, we were concerned with establishing whether the social and interpersonal improvements found at the end of 36 months had been maintained and whether additional gains in the area of vocational achievement had been made in either group. We also looked at continuing use of medical and psychiatric services, including emergency room visits, length of hospitalization, outpatient psychiatric care, community support, use of medication and psychological therapies, and overall symptom status. This article reports on these long-term outcomes for patients who participated in the original trial.

Method

The characteristics of the subjects, the methodology of the original trial, and the details of treatment have been described (15, 17). Both groups had access to inpatient treatment for acute crises if recommended by the primary psychiatrist. At the end of 18 months, the mentalization-based treatment by partial hospitalization patients were offered twice-weekly outpatient mentalizing group psychotherapy for a further 18 months, whereas the treatment as usual group continued with general psychiatric care with psychotherapy but not mentalization-based treatment if recommended by the consultant psychiatrist.

Mentalization-based treatment by partial hospitalization consists of 18-month individual and group psychotherapy in a partial hospital setting offered within a structured and integrated program provided by a supervised team. Expressive therapy using art and writing groups is included. Crises are managed within the team; medication is prescribed according to protocol by a psychiatrist working in the therapy program. The understanding of behavior in terms of underlying mental states forms a common thread running across all aspects of treatment. The focus of therapy is on the patient's moment-to-moment state of mind. The patient and therapist collaboratively try to generate alternative perspectives to the patient's subjective experience of himself or herself and others by moving from validating and supportive interventions to exploring the therapy relationship itself as it suggests alternative understanding. This psychodynamic therapy is manualized (17) and in many respects overlaps with transference-focused psychotherapy (18).

Treatment as usual consists of general psychiatric outpatient care with medication prescribed by the consultant psychiatrist, community support from mental health nurses, and periods of partial hospital and inpatient treatment as necessary but no specialist psychotherapy.

We initially reported conservatively on all patients randomly assigned to the mentalization-based treatment by partial hospitalization/group therapy condition regardless of their duration of treatment at 36 months, including dropouts (16). In the current study, we followed up all 41 patients 8 years after random assign-

ment (5 years after they had ceased all mentalization-based treatment). Contact was made by letter, through their general practitioner, and by telephone. Written informed consent was obtained in person or by letter after the follow-up study had been fully explained according to the requirements of the local research ethics committee. Medical and psychiatric records were obtained for all 41 patients and relevant information extracted. The health service in the United Kingdom requires patients to have treatment in their local area. Tertiary care medical records enable tracing and estimation of health care use over long periods.

The patients in the study group were interviewed by research psychologists who remained blind to original group allocation. One patient in the treatment as usual group had committed suicide. Five patients (three in treatment as usual and two in mentalization-based treatment by partial hospitalization) refused a personal interview, citing schedule or travel problems. The two mentalization-based treatment by partial hospitalization patients accepted a telephone interview.

Assessment

The primary outcome measure was the number of suicide attempts over the whole of the 5-year postdischarge follow-up period. Associated outcomes were service use, including emergency room visits; the length and frequency of hospitalization; continuing outpatient psychiatric care; and use of medication, psychological therapies, and community support.

Secondary outcomes were 1) symptom status as assessed at a follow-up interview using the Zanarini Rating Scale for DSM-IV borderline personality disorder (19) and 2) global functioning as measured by the Global Assessment of Functioning Scale (GAF), which has been found to show less improvement in naturalistic follow-along studies than diagnostic symptom profiles (20).

At 6-month intervals after 18 months of mentalization-based treatment by partial hospitalization, we assessed treatment profiles (emergency room visits, hospitalization, psychiatric outpatients, community support, psychotherapy, medication) and suicidality and self-harm using criteria defined in the original trial for each patient by interview and scrutiny of medical records. We also collected information twice yearly concerning vocational status, calculating the number of 6-month periods in which the patient was employed or attended an educational program for more than 3 months. Patient recall for self-harm was unreliable and could not be independently corroborated from medical records and so is not reported. However, we consider the frequency of emergency room visits to be a reasonable proxy of severe self-harm in this population.

The reliability of information gained from medical records was assessed on a random subset of notes (45%), which were independently coded by two researchers. A similar proportion of recorded interviews was assessed for interrater agreement. Although interviews could be conducted blind, data extraction from the medical notes could not be performed without knowledge of treatment allocation. To reduce bias, all pertinent data (e.g., suicide attempts, hospitalization, emergency room visits, vocation) were cross-checked with other sources of information (e.g., emergency room, general practitioner, education institution records). Inter-coder agreement was in excess of 90% for almost all variables used (median kappa=0.90, range=0.77–1.00, for each 6-month period). Final GAF scores were assigned independently by two blinded judges on the basis of current case notes and interview information; interrater reliability was 0.72. Aggregate scores were used in the analysis.

The primary outcome measure of suicide had an extremely skewed distribution, and so nonparametric Mann-Whitney statistics were applied to frequency data. We used the Mann-Whitney test or analysis of variance depending on the distribution for the other variables. Multivariate analysis of variance was used to

Patient Perspectives

A 24-year-old female patient was referred from forensic services after her arrest for setting fire to her university dormitories. She had a history of recent suicide attempts and regularly burned herself with cigarettes and a hot iron. Feelings of rejection in her current relationship with her partner could have triggered serious self-harm. She was admitted to the mentalization-based treatment by partial hospitalization program and offered individual (one session per week) and group psychotherapy (three sessions per week) with the addition of art therapy (two sessions per week) within the expressive therapy program. The program was organized over 5 days and amounted to 9 hours of therapy per week with 3-monthly review of her antipsychotic and antidepressant medication. In individual sessions, treatment initially focused on clarifying her own feelings and others' experience of her. The eventual focus was on how her experiences of self-doubt and emotional turbulence led to a sense of fragmentation that was controlled only by experiences of intense physical pain. The individual therapist identified these processes while focusing on the way she represented her own mental states and those of others with whom she interacted. Gradually this was explored within the relationship with the therapist: "it never occurred to me that what I did had an effect on anyone else." In groups, the patient was frequently challenged about the effect of her behavior on other group members. She frequently threatened to leave the group. The individual and group therapist collaborated in helping to maintain her attendance in treatment. In art therapy, she was encouraged to express her inner states in her painting and to explain her pictures to others and to consider others' understanding of and reactions to them.

During the treatment, she terminated her relationship with her abusive partner and stopped her medication. She reentered college and continued with mentalization-based treatment with group therapy. At the end of 36 months, she was discharged and a year later joined training courses for professionals wishing to learn more about mentalization-based treatment.

A 28-year-old female patient was randomly assigned to treatment as usual and returned to treatment with her referring outpatient psychiatrist and to the community support team. A mental health nurse and psychologist agreed to target her self-harm and social problems using problem-solving techniques and support in a crisis. She saw the psychologist weekly for 3 months. Her self-harm improved initially, but a serious suicide attempt led to inpatient admission under the care of a different psychiatrist, who changed the patient's antidepressant medication to antipsychotic medication and added a mood stabilizer with occasional use of benzodiazepines for anxiety. At discharge, the patient made a formal complaint against the hospital for failure to ensure appropriate discharge planning. She was transferred to partial hospital care, where she improved. When her psychiatrist tried to reduce her benzodiazepines, her self-harm became more frequent. Following discharge from partial hospitalization, she attended psychiatric outpatient care for a further 6 months and refused to see the psychologist but made a good relationship with the support nurse, who met with her regularly at home and was available in a crisis. She continued to be seen as a psychiatric outpatient for a further year. Her main complaint at interview was that "no one seemed to understand what I needed."

contrast the two groups on the Zanarini Rating Scale for Borderline Personality Disorder. For service use (outpatient psychiatry, community support, and psychotherapy), we computed the percentage of available services used for each patient for the year before random assignment and during subsequent blocks of time (mentalization-based treatment by partial hospitalization, 18 months), mentalization-based treatment by group (18 months), and postdischarge (0–18 months, 19–36 months, 37–60 months). For the same periods, we also computed the proportion of each group who were hospitalized, made suicide attempts, were employed or in education, attended the emergency room, and were taking three or more classes of medication. For each time block, the proportions were contrasted using chi-square statistics.

Results

Means and standard deviations of primary and secondary outcomes for mentalization-based treatment and treatment as usual groups are shown in Table 1 covering

the 5-year postdischarge period together with significant statistics and effect sizes contrasting the two groups. For frequency, data effect sizes are stated as numbers needed to treat (Newcombe-Wilson 95% confidence interval [CI]).

Overall, 46% of the patients made at least one suicide attempt (one successfully), but only 23% did so in the mentalization-based treatment group, contrasted with 74% of the treatment as usual group. There was a significant difference on the Mann-Whitney U test in the total number of suicide attempts over the follow-up period. Figure 1 shows the percentage of each group that made a suicide attempt during each block of time. Significant differences between the groups were apparent during the mentalization-based treatment group therapy period and remained significant in all three postdischarge periods.

Table 1 shows that the mean number of emergency room visits and hospital days highly significantly favored

TABLE 1. Effect Sizes for Primary and Secondary Outcomes for Mentalization-Based Treatment by Partial Hospitalization/Group Therapy and Treatment as Usual Groups Over 5 Years Postdischarge

Measure	Mentalization-Based Treatment by Partial Hospitalization/Group Therapy (N=22)		Treatment as Usual (N=19)		Analysis			Effect Size ^a	
	Mean	SD	Mean	SD	Test	df	p	d	95% CI
Suicide attempts									
Total number	0.05	0.9	0.52	0.48	U=73, z=3.9		0.00004	1.4	1.3 to 1.5
Any attempt	N	%	N	%	Test	df	p	d	95% CI
	5	23	14	74	$\chi^2=8.7$	1	0.003	2.0	1.4 to 4.9
Zanarini Rating Scale for Borderline Personality Disorder ^b									
Positive criteria	3	14	13	87	$\chi^2=16.5$	1	0.000004	1.4	1.2 to 2.4
	Mean	SD	Mean	SD	Test	df	p	d	95% CI
Total	5.5	5.2	15.1	5.3	F=29.7	1, 35	0.000004	1.80	0.14 to 3.50
Affect	1.6	2.0	3.7	2.0	F=9.7	1, 35	0.004	1.10	0.41 to 1.70
Cognitive	1.1	1.4	2.5	2.0	F=6.9	1, 35	0.02	0.84	0.30 to 1.40
Impulsivity	1.6	1.8	4.1	2.3	F=13.9	1, 35	0.001	1.20	0.59 to 1.90
Interpersonal	1.5	1.7	4.7	2.3	F=23.2	1, 35	0.00003	1.6	1.0 to 2.3
GAF score ^c	58.3	10.5	51.8	5.7	F=5.4	1, 35	0.03	0.75	-1.90 to 3.40
	N	%	N	%	Test	df	p	d	95% CI
GAF score >61	10	46	2	11	$\chi^2=6.5$	1	0.02	3	2 to 12
	Mean	SD	Mean	SD	Test	df	p	d	95% CI
Number of days of hospitalization ^c	0.27	0.71	6.2	5.6	U=25.5, z=5.1		0.00000002	1.50	0.36 to 2.70
Number of emergency room visits ^c	0.77	1.10	6.4	5.7	U=66.0, z=3.9		0.00003	1.40	0.21 to 2.63
Number of years of employment ^c	3.2	2.3	1.2	1.9	F=8.9	1, 35	0.005	0.94	0.29 to 1.60
Number of years of further treatment ^c									
Further psychiatric outpatient treatment	2.0	1.9	3.6	1.5	F=8.5	1, 35	0.006	0.93	-4.00 to 1.50
Further therapy 36 months postintake	0.48	1.10	0.55	0.83	F=0.6	1, 35	n.s.	0.07	-0.23 to 0.37
Further assertive outreach treatment	0.39	0.51	2.7	1.8	U=33.5, z=4.68		0.0000002	1.8	1.4 to 2.2
Medication (years) ^c									
Antidepressants	1.1	1.8	3.3	2.3	F=11.6	1, 35	0.002	1.10	0.45 to 1.70
Antipsychotics	0.16	0.28	3.1	2.1	U=9.0, z=5.4		0.000000005	2.04	1.60 to 2.50
Mood stabilizers	0.11	0.26	1.8	2.1	U=105.0, z=3.2		0.001	1.17	0.73 to 1.60
Three or more drugs (including hypnotics)	0.02	0.11	1.9	1.9	U=58.5, z=4.6		0.0000009	1.45	1.10 to 1.80

^a For frequency variables, data effect sizes are stated as numbers needed to treat with Newcombe-Wilson 95% confidence intervals.

^b Number for treatment as usual=15.

^c Number for treatment as usual=18.

the mentalization-based treatment group, as did the continuing treatment profile. Figure 1 shows the percentage of patients in each group who made an emergency room visit and were hospitalized at least once during the study periods. Emergency room visits were significantly reduced in all periods of treatment and postdischarge. The percent hospitalized was significantly lower during the last two postdischarge periods.

During mentalization-based treatment group therapy, all of the experimental group but only 31% of the treatment as usual group received therapy ($\chi^2=21$, $df=1$, $p=0.0000005$). Over the 5-year postdischarge period, both groups received around 6 months of psychological therapy (n.s.). For all other treatments, the treatment as usual group received significantly more input postdischarge—

3.6 years of psychiatric outpatient treatment and 2.7 years of assertive community support, compared with 2 years and 5 months, respectively, for the mentalization-based treatment group. The mean percent of available services used throughout the period of the study is shown in Figure 1. The differences favored the treatment as usual group only in the initial treatment period (mentalization-based treatment by partial hospitalization) and were significantly less for the mentalization-based treatment group for all three postdischarge periods.

Differences were also marked in terms of medication (Table 1). The treatment as usual group had an average of over 3 years taking antipsychotic medication, whereas the mentalization-based treatment group had less than 2 months. Somewhat smaller but still substantial differ-

FIGURE 1. Pretreatment Levels and Outcomes for Mentalization-Based Treatment by Partial Hospital/Group and Treatment as Usual Groups Over 8 Years Postrandomization



^a Services included outpatient psychiatry, community support, and psychotherapy. Probabilities refer to chi-square statistics.

* $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$.

ences were apparent in antidepressant and mood stabilizer use. The treatment as usual group spent nearly 2 years taking three or more psychoactive medications, compared to an average of 2 months for the mentalization-based treatment group. Figure 1 shows that around 50% of the treatment as usual patients but none of the mentalization-based treatment group were taking three or more classes of psychoactive medication during mentalization-based treatment group therapy and the three post-discharge periods.

At the end of the follow-up period, 13% of the mentalization-based treatment patients met diagnostic criteria for borderline personality disorder, compared with 87% of the treatment as usual group (Table 1). The contrast between mean total scores for the Zanarini Rating Scale for Borderline Personality Disorder yielded a large effect size favoring the mentalization-based treatment group, albeit with a wide confidence interval. Multivariate analysis of variance across the four symptom clusters also reflected the better outcome for the mentalization-based treatment

group (Wilks's $\lambda=0.55$, $F=6.4$, $df=4$, 32 , $p=0.001$). The largest differences favoring mentalization-based treatment were in terms of impulsivity and interpersonal functioning. Table 1 shows there was over a 6-point difference in the GAF scores between the two groups, yielding a clinically significant moderate effect size of 0.8 (95% CI=-1.9 to 3.4). Forty-six percent of the mentalization-based treatment group compared to 11% of the treatment as usual group had GAF scores above 60. Of importance, vocational status favored the mentalization-based treatment group, who were employed for nearly three times as long as the treatment as usual group. Figure 1 shows a gradual increase in the percent of mentalization-based treatment patients in employment or education in the three postdischarge periods.

Discussion

The mentalization-based treatment by partial hospitalization/group therapy group continued to do well 5 years after all mentalization-based treatment had ceased. The beneficial effect found at the end of mentalization-based treatment group therapy for borderline personality disorder is maintained for a long period, with differences found in suicide attempts, service use, global function, and Zanarini Rating Scale for Borderline Personality Disorder scores at 5 years postdischarge. It is consistent with the possible rehabilitative effects that we observed during the mentalization-based treatment group therapy period. This is encouraging because positive effects of treatment normally tend to diminish over time. The treatment as usual group received more treatment over time than the mentalization-based treatment group, perhaps because they continued to have more symptoms. However, in both groups, GAF scores continue to indicate deficits, with some patients continuing to show moderate difficulties in social and occupational functioning. Nevertheless, when compared to the treatment as usual group, mentalization-based treatment by partial hospital/group therapy patients were more likely to be functioning reasonably well with some meaningful relationships as defined by a score higher than 60.

More striking than how well the mentalization-based treatment group did was how badly the treatment as usual group managed within services despite significant input. They look little better on many indicators than they did at 36 months after recruitment to the study. A few patients in the mentalization-based treatment group had made at least one suicide attempt during the postdischarge period, but this was almost 10 times more common in the treatment as usual group. Associated with this were more emergency room visits and greater use of polypharmacy. However, although the number of hospital days was greater for the treatment as usual group than the mentalization-based treatment group, the percentage of patients admitted to the hospital over the postdischarge period

was small (25%–33%). This pattern of results suggests not that treatment as usual is necessarily ineffective in its components but that the package or organization is not facilitating possible natural recovery.

Naturalistic follow-up studies suggest spontaneous remission of impulsive symptoms within 2–4 years with apparently less treatment (21, 22). In line with these findings, all patients showed improvement, although not as much in terms of suicide attempts as might be expected. The lower level of improvement observed in this population may represent a more chronically ill group. Most patients had a median time in specialist services at entry to the trial of 6 years. Although this study does not indicate the untreated course of the disorder, the results suggest that quantity of treatment may not be a good indicator of improvement and may even prevent patients from taking advantage of felicitous social and interpersonal events (23). It is possible that treatment as usual inadvertently interfered with patient improvement as well as mentalization-based treatment accelerating recovery.

There is an anomaly in the results in that there is a marked difference between the size of the effects as measured by the Zanarini Rating Scale for Borderline Personality Disorder and the GAF in terms of social and interpersonal function. One possible explanation for this is that the scales offer a slightly different metric to different aspects of interpersonal function. In the GAF, suicidal preoccupation and actual attempts have a large loading, and even presence of suicidal thoughts reduces the score substantially. This was the case for a small number of patients in the mentalization-based treatment group and accounts for their larger variance on GAF scores. In contrast, the interpersonal subscale of the Zanarini Rating Scale for Borderline Personality Disorder covers two symptoms in the interpersonal realm of borderline personality disorder, namely, intense unstable relationships and frantic efforts to avoid abandonment, that showed marked improvement in the mentalization-based treatment group. A GAF of greater than 60 clearly marks a change back to improved function, and more patients in the mentalization-based treatment group achieved scores above this level. A strong correlate of improvement in the mentalization-based treatment group is vocational status. It is unclear whether this is a cause or consequence of improvement. It is likely that symptomatic improvement and vocational activity represent a virtuous cycle. Although we have no evidence to this effect, we suggest that mentalization-based treatment may be specifically helpful in improving patient ability to manage social situations by enabling individuals to distance themselves from the interpersonal pressures of the work situation, anticipate other people's thoughts and feelings, and be able to understand their own reactions without overactivation of their attachment systems (24, 25).

The strengths of this study lie in the presence of a long-term control group, in the reliability of care records, and

in our data collection for suicide attempts, which used the same rigorous criteria as at the outset of the trial. Other follow-up studies have been confounded by a lack of controls or treatment as usual patients being taken in to the experimental treatment at the end of the treatment phase. However, the long-term follow-up of a small group and allegiance effects, despite attempts being made to blind the data collection, limit the conclusions. In addition, some of the measures we used at the outset of the trial were not repeated in this follow-up. We considered the Zanarini Rating Scale for Borderline Personality Disorder to be a more useful outcome measure that would reflect the current state of the patients better than self-report questionnaire methods. Finally, the original mentalization-based treatment by partial hospitalization intervention contained a number of components in addition to psychological therapy. It is therefore unclear whether psychodynamic therapy was the essential component. In order for mentalization-based treatment to be accepted as an evidence-based treatment for borderline personality disorder, larger trials using core components of the intervention are necessary. These are now being undertaken. Although this study demonstrates that borderline patients improve in a number of domains after mentalization-based treatment and that those gains are maintained over time, global function remains somewhat impaired. This may reflect too great a focus during treatment on symptomatic problems at the expense of concentration on improving general social adaptation.

Received April 15, 2007; revised Sept. 11, 2007; accepted Dec. 14, 2007 (doi: 10.1176/appi.ajp.2007.07040636). From St. Ann's Hospital, Barnet, Enfield, and Haringey Mental Health Trust, London; and the Department of Clinical Health Psychology, University College London, London. Address correspondence and reprint requests to Dr. Bateman, Halliwick Unit, St. Ann's Hospital, St. Ann's Rd., London, U.K. N15 3TH; anthony@abate.org.uk (e-mail).

All authors report no competing interests.

The authors are in receipt of a grant from the Borderline Personality Disorder Research Foundation. Supported by a grant from Barnet, Enfield, and Haringey Mental Health NHS Trust.

The authors thank Carmen Chan, Joanna Jackson, and Bethany Taylor for their assistance with this project.

References

- Brazier J, Tumor I, Holmes M, Ferriter M, Parry G, Dent-Brown K, Paisley S: Psychological therapies including dialectical behaviour therapy for borderline personality disorder: a systematic review and preliminary economic evaluation. *Health Technology Assessment* 2006; 10:iii, ix–xii, 1–117
- Linehan MM, Comtois KA, Murray AM, Brown MZ, Gallop RJ, Heard HL, Korslund KE, Tutek DA, Reynolds SK, Lindenboim N: Two-year randomized controlled trial and follow-up of dialectical behavior therapy vs therapy by experts for suicidal behaviors and borderline personality disorder. *Arch Gen Psychiatry* 2006; 63:757–766
- Koons CR, Robins CJ, Tweed JL, Lynch TR, Gonzalez AM: Efficacy of DBT in women veterans with BPD. *Behav Ther* 2001; 32: 371–390
- Turner RM: Naturalistic evaluation of DBT oriented treatment for BPD. *Cogn Behaviour Practice* 2000; 7:413–419
- Giesen-Bloo J, van Dyck R, Spinhoven P, van Tilburg W, Dirksen C, van Asselt T, Kremers I, Nadort M, Arntz A: Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transference-focused psychotherapy. *Arch Gen Psychiatry* 2006; 63:649–658
- Davidson K, Norrie J, Tyrer P, Gumley A, Tata P, Murray H, Palmer S: The effectiveness of cognitive behavior therapy for borderline personality disorder: results from the Borderline Personality Disorder Study of Cognitive Therapy (BOSCO) trial. *J Personal Disord* 2006; 20:450–465
- Clarkin JF, Levy KN, Lenzenweger MF, Kernberg OF: Evaluating three treatments for borderline personality disorder: a multi-wave study. *Am J Psychiatry* 2007; 164:922–928
- Zanarini MC, Frankenburg FR, Hennen J, Reich DB, Silk K: Psychosocial functioning of borderline patients and axis II comparison subjects followed prospectively for six years. *J Personal Disord* 2005; 19:19–29
- Skodol AE, Pagano ME, Bender DS, Shea MT, Gunderson JG, Yen S, Stout RL, Morey LC, Sanislow CA, Grilo CM, Zanarini MC, McGlashan TH: Stability of functional impairment in patients with schizotypal, borderline, avoidant, or obsessive-compulsive personality disorder over two years. *Psychol Med* 2005; 35:443–451
- Zanarini MC, Frankenburg FR, Reich DB, Silk KR, Hudson JI, McSweeney LB: The subsyndromal phenomenology of borderline personality disorder: a 10-year follow-up study. *Am J Psychiatry* 2007; 164:929–935
- Linehan MM, Heard HL, Armstrong HE: Naturalistic follow-up of a behavioral treatment for chronically parasuicidal borderline patients. *Arch Gen Psychiatry* 1993; 50:971–974
- van den Bosch LM, Koeter MW, Stijnen T, Verheul R, van den Brink W: Sustained efficacy of dialectical behaviour therapy for borderline personality disorder. *Behav Res Ther* 2005; 43: 1231–1241
- Stevenson J, Meares R, D'Angelo R: Five-year outcome of outpatient psychotherapy with borderline patients. *Psychol Med* 2005; 35:79–87
- Chiesa M, Fonagy P, Holmes J: Six-year follow-up of three treatment programs to personality disorder. *J Personal Disord* 2006; 20:493–509
- Bateman A, Fonagy P: Effectiveness of partial hospitalization in the treatment of borderline personality disorder: a randomized controlled trial. *Am J Psychiatry* 1999; 156:1563–1569
- Bateman A, Fonagy P: Treatment of borderline personality disorder with psychoanalytically oriented partial hospitalization: an 18-month follow-up. *Am J Psychiatry* 2001; 158:36–42
- Bateman AW, Fonagy P: *Mentalization Based Treatment for Borderline Personality Disorder: A Practical Guide*. Oxford, UK, Oxford University Press, 2006
- Kernberg OF, Clarkin JF, Yeomans FE: *A Primer of Transference Focused Psychotherapy for the Borderline Patient*. New York, Jason Aronson, 2002
- Zanarini MC: Zanarini Rating Scale for Borderline Personality Disorder (ZAN-BPD): a continuous measure of DSM-IV borderline psychopathology. *J Personal Disord* 2003; 17:233–242
- Gunderson JG, Daversa MT, Grilo CM, McGlashan TH, Zanarini MC, Shea MT, Skodol AE, Yen S, Sanislow CA, Bender DS, Dyck IR, Morey LC, Stout RL: Predictors of 2-year outcome for patients with borderline personality disorder. *Am J Psychiatry* 2006; 163:822–826
- McGlashan TH, Grilo CM, Sanislow CA, Ralevski E, Morey LC, Gunderson JG, Skodol AE, Shea MT, Zanarini MC, Bender D, Stout RL, Yen S, Pagano M: Two-year prevalence and stability of individual DSM-IV criteria for schizotypal, borderline, avoidant,

- and obsessive-compulsive personality disorders: toward a hybrid model of axis II disorders. *Am J Psychiatry* 2005; 162:883–889
22. Skodol AE, Gunderson JG, Shea MT, McGlashan TH, Morey LC, Sanislow CA, Bender DS, Grilo CM, Zanarini MC, Yen S, Pagano ME, Stout RL: The Collaborative Longitudinal Personality Disorders Study (CLPS): overview and implications. *J Personal Disord* 2005; 19:487–504
23. Fonagy P, Bateman A: Progress in the treatment of borderline personality disorder. *Br J Psychiatry* 2006; 188:1–3
24. Levy KN, Meehan KB, Kelly KM, Reynoso JS, Weber M, Clarkin JF, Kernberg OF: Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. *J Consult Clin Psychol* 2006; 74:1027–1040
25. Fonagy P, Bateman AW: Mechanisms of change in mentalization-based treatment of BPD. *J Clin Psychol* 2006; 62:411–430