

### Scientifically Unfounded Claims in Diagnosing and Treating Patients

TO THE EDITOR: We greatly appreciated the thoughtful book review by Andrew F. Leuchter, M.D. (1), published in the May 2009 issue of the *Journal*, on Daniel Amen's *Healing the Hardware of the Soul: Enhance Your Brain to Improve Your Work, Love, and Spiritual Life* (2). Dr. Amen claims that numerous psychiatric illnesses can be diagnosed and treatments prescribed based on resting single photon emission computerized tomography (SPECT) images. Dr. Leuchter correctly points out the absence of empirical data to support the claims of Dr. Amen. Several years ago, following conversations with Dr. Amen on how to address such concerns, the Brain Imaging Council of the Society of Nuclear Medicine offered Dr. Amen the opportunity to submit his analyses of a blinded set of SPECT scans (to have been prepared by the Brain Imaging Council) to determine how effective his technique is at correctly diagnosing subjects. Although this proposed study could have provided support for his approach, the offer was declined. Nevertheless, for more than two decades, Dr. Amen has persisted in using scientifically unfounded claims to diagnose and treat patients (over 45,000 by his own count).

There are several dangers to patients that can accrue from this approach: 1) patients (including children) are administered a radioactive isotope without sound clinical rationale; 2) patients pursue treatments contingent upon an interpretation of a SPECT image that lacks empirical support; and 3) based on a presumed diagnosis provided by Dr. Amen's clinics, patients are guided toward treatment that may detract them from clinically sound treatments.

Just as serious is the danger to our field. It is likely that, within the next decade, Dr. Amen's claims will be realized in that psychiatrists will enjoy the ability to diagnose and prescribe treatments based, in part, upon neuroimaging findings. Unfortunately, if previously led astray by unsupported claims, patients and their doctors may be less inclined to utilize scientifically proven approaches once these are shown in the peer-reviewed literature to be effective.

It is therefore incumbent upon all of us to monitor and regulate our field. We encourage physicians to remain vigilant of unproven approaches practiced by our peers and to immediately report these trespasses to their state medical boards.

#### References

1. Leuchter AF: Healing the hardware of the soul: enhance your brain to improve your work, love, and spiritual life, by Daniel Amen (book review). *Am J Psychiatry* 2009; 166:625
2. Amen D: *Healing the Hardware of the Soul: Enhance Your Brain to Improve Your Work, Love, and Spiritual Life*. New York, Simon and Schuster, 2008

BRYON ADINOFF, M.D.  
MICHAEL DEVOUS, Ph.D.  
Dallas, Tex.

*Dr. Adinoff has received grant/research support from the Department of Veterans Affairs, the National Institute on Alcohol Abuse and Alcoholism, and the National Institute on Drug Abuse; he has served as a consultant to GlaxoSmithKline, the Hershowe Law Firm, Phillips Lytle (for GlaxoSmithKline), Shook, Hardy and Bacon, Simon Pissante, and Teva Pharma-*

*ceutical Industries; and he has received honoraria from the American Academy of Addiction Psychiatry, the Medical University of South Carolina, and the University of New Mexico. Dr. Devous has received research support from Alseres and AVID Radiopharmaceuticals, and he has served on the scientific advisory board of AVID Radiopharmaceuticals.*

*This letter (doi: 10.1176/appi.ajp.2010.10020157) was accepted for publication in March 2010.*

### Psychosis Associated With Medical Marijuana: Risk vs. Benefits of Medicinal Cannabis Use

TO THE EDITOR: Over the past 15 years, it has become increasingly evident that cannabis use carries an increased risk for the development of psychosis (1, 2). At the same time, medicinal cannabis (medical marijuana) has been legalized in many states, with minimal restrictions on prescribing indications. The present case illustrates the evolution of a psychotic disorder, in the setting of medicinal cannabis use, in a young man at high risk for psychosis.

"Mr. Z" was a 24-year-old man who was first hospitalized for insomnia, irritability, and aggressiveness 2 years after military service. On admission, he displayed heightened religiosity and mild suspiciousness. Urine toxicology screening revealed cannabinoids, supporting the patient's endorsed semi-daily cannabis use via water pipe for the past 18 months, without other substance abuse. He was started on quetiapine (100 mg/day), with rapid resolution of symptoms, and discharged after 10 days.

The patient subsequently discontinued quetiapine and was lost to follow-up. Four months later, he presented to a marijuana clinic complaining of chronic pain, insomnia, and anxiety and was given a diagnosis of posttraumatic stress disorder (PTSD) and pain, along with a medical recommendation for cannabis. No psychotic symptoms were elicited. He later explained that he switched from "street" marijuana to medical marijuana in order to obtain a more potent product as well as to avoid illegal activity and getting "ripped off" by drug dealers. He also increased the frequency of his daily use from approximately once to twice daily.

Six months later, Mr. Z was rehospitalized with new-onset auditory hallucinations (multiple voices speaking to each other and urging violence) and delusions (believing that people were tampering with his windows and eavesdropping on his conversations and that he was Jesus Christ). Aripiprazole (15 mg/day) was prescribed, with gradual symptomatic improvement, and then tapered to a lower dose (7.5 mg/day) due to tremor. The patient reported that he believed smoking cannabis helped his chronic pain but that it worsened his psychotic symptoms, such that he wanted help to stop smoking the drug. After 4 weeks, he was discharged to residential substance abuse treatment with only mild, residual psychotic symptoms and a discharge diagnosis of psychotic disorder not otherwise specified, PTSD, and cannabis dependence. At a 3-month follow-up evaluation, while still taking aripiprazole, Mr. Z remained off cannabis and free of psychotic symptoms.

Although cannabis may have some health benefits, it also has a variety of adverse effects, including psychosis, especially among those at high risk (1–3). The patient in the present case was at high risk for psychosis based on attenuated symp-