

Genome-Wide Association Study Meta-Analysis of the Alcohol Use Disorders Identification Test (AUDIT) in Two Population-Based Cohorts

Sandra Sanchez-Roige, Ph.D., Abraham A. Palmer, Ph.D., Pierre Fontanillas, Ph.D., Sarah L. Elson, Ph.D., the 23andMe Research Team, the Substance Use Disorder Working Group of the Psychiatric Genomics Consortium, Mark J. Adams, Ph.D., David M. Howard, Ph.D., Howard J. Edenberg, Ph.D., Gail Davies, Ph.D., Richard C. Crist, Ph.D., Ian J. Deary, Ph.D., M.R.C.Psych., Andrew M. McIntosh, M.D., Toni-Kim Clarke, Ph.D.

Objective: Alcohol use disorders are common conditions that have enormous social and economic consequences. Genome-wide association analyses were performed to identify genetic variants associated with a proxy measure of alcohol consumption and alcohol misuse and to explore the shared genetic basis between these measures and other substance use, psychiatric, and behavioral traits.

Method: This study used quantitative measures from the Alcohol Use Disorders Identification Test (AUDIT) from two population-based cohorts of European ancestry (UK Biobank [N=121,604] and 23andMe [N=20,328]) and performed a genome-wide association study (GWAS) meta-analysis. Two additional GWAS analyses were performed, a GWAS for AUDIT scores on items 1–3, which focus on consumption (AUDIT-C), and for scores on items 4–10, which focus on the problematic consequences of drinking (AUDIT-P).

Results: The GWAS meta-analysis of AUDIT total score identified 10 associated risk loci. Novel associations localized to genes including *JCAD* and *SLC39A13*; this study also replicated previously identified signals in the genes *ADH1B*, *ADH1C*, *KLB*, and *GCKR*. The dimensions of AUDIT showed positive

genetic correlations with alcohol consumption ($r_g=0.76-0.92$) and DSM-IV alcohol dependence ($r_g=0.33-0.63$). AUDIT-P and AUDIT-C scores showed significantly different patterns of association across a number of traits, including psychiatric disorders. AUDIT-P score was significantly positively genetically correlated with schizophrenia ($r_g=0.22$), major depressive disorder ($r_g=0.26$), and attention deficit hyperactivity disorder ($r_g=0.23$), whereas AUDIT-C score was significantly negatively genetically correlated with major depressive disorder ($r_g=-0.24$) and ADHD ($r_g=-0.10$). This study also used the AUDIT data in the UK Biobank to identify thresholds for dichotomizing AUDIT total score that optimize genetic correlations with DSM-IV alcohol dependence. Coding individuals with AUDIT total scores ≤ 4 as control subjects and those with scores ≥ 12 as case subjects produced a significant high genetic correlation with DSM-IV alcohol dependence ($r_g=0.82$) while retaining most subjects.

Conclusions: AUDIT scores ascertained in population-based cohorts can be used to explore the genetic basis of both alcohol consumption and alcohol use disorders.

Am J Psychiatry 2019; 176:107–118; doi: 10.1176/appi.ajp.2018.18040369

Alcohol use disorders are modestly heritable, with twin studies demonstrating that approximately 50% of the phenotypic variance is attributed to genetic factors (1, 2). To date, genetic studies of alcohol use disorders have identified genes that influence pharmacokinetic factors (e.g., *ADH1B*, *ADH1C*, *ALDH2*) (3–8) but none that influence pharmacodynamic factors. The difficulty of assembling large, carefully diagnosed cohorts of individuals with alcohol use disorders has stimulated additional studies of nonclinical phenotypes, such as alcohol consumption, in populations not ascertained for alcohol dependence. This approach has allowed for the relatively rapid collection of much larger sample sizes (e.g., >100,000

individuals) and has identified numerous loci associated with both pharmacokinetic and pharmacodynamic factors that influence alcohol consumption, including *ADH1B*/*ADH1C*/*ADH5* (9–11), *KLB* (encoding β -klotho) (9, 11, 12), and *GCKR*, encoding the glucokinase regulatory protein (9, 11). However, the genetic overlap between alcohol consumption (in units per week) and diagnosed DSM-IV alcohol dependence is moderate ($r_g=0.38$) (13), which reinforces the notion that alcohol consumption cannot be used as a surrogate for alcohol use disorders.

The Alcohol Use Disorders Identification Test (AUDIT) is a screening tool designed to identify past-year hazardous

See related features: Editorials by Dr. Brady (p. 87) and Dr. Pollock and Dr. Lossie (p. 92) and **AJP Audio** (online)

alcohol use (14). The test consists of 10 items across three dimensions—those pertaining to alcohol consumption (items 1–3, often termed AUDIT-C), dependence symptoms (items 4–6), and problematic or hazardous alcohol use (items 7–10). When the AUDIT was developed, a total score ≥ 8 was proposed to be indicative of hazardous alcohol use (14) and a score ≥ 20 consistent with a diagnosis of alcohol dependence (15). However, there is no clear consensus on the threshold for alcohol dependence, and subsequent studies have suggested that additional factors, including sex and cultural and social contexts, should be considered when deriving thresholds for alcohol dependence (reviewed in Table S1 in the online supplement).

A recent population-based GWAS of AUDIT scores in 20,328 research participants from the genetics company 23andMe identified a locus near the gene *ADH1C* (rs141973904; $p=4.4 \times 10^{-7}$) (10) that was nominally associated with AUDIT total score. AUDIT scores among 23andMe research participants were low and predominantly driven by alcohol consumption (AUDIT-C score). The genetic correlation between AUDIT total score from the 23andMe sample and alcohol consumption was much stronger ($r_g=0.89$, $p=9.01 \times 10^{-10}$) than the genetic correlation between AUDIT total score and alcohol dependence ($r_g=0.08$; $p=0.65$) (13).

In this study, we performed a GWAS meta-analysis using the UK Biobank cohort ($N=121,604$) and the previously published 23andMe cohort ($N=20,328$) (10), yielding the largest GWAS meta-analysis of AUDIT total score to date ($N=141,932$). Using only the UK Biobank cohort, we also sought to determine whether the alcohol consumption component of the AUDIT had a genetic architecture distinct from the dependence and hazardous-use components by performing GWASs of consumption (items 1–3, AUDIT-C) and problems (items 4–10, AUDIT-P). Linkage disequilibrium score regression (16) was used to calculate genetic correlations between AUDIT measures and other substance use, psychiatric, and behavioral traits. We also calculated genetic correlations with obesity and blood lipid traits, as these have previously been shown to be associated with alcohol consumption (9, 10). We hypothesized that AUDIT-P score would correlate more strongly with measures of hazardous substance use, including alcohol dependence, and other psychiatric conditions. Finally, to determine the thresholds for dichotomizing AUDIT total score that would most closely approximate alcohol dependence, we used continuous AUDIT total score to categorize participants as case and control subjects using different thresholds, performed GWAS on each, and calculated the genetic correlation with DSM-IV alcohol dependence (13).

METHOD

UK Biobank Sample

The UK Biobank (www.ukbiobank.ac.uk) is a population-based sample of 502,629 individuals who were recruited from 22 assessment centers across the United Kingdom from 2006 to 2010 (17). A total of 157,366 individuals completed a

mental health questionnaire as part of an online follow-up over a 1-year period in 2017. The AUDIT was administered to assess alcohol use over the past year, using gating logic (see Figure S1 in the online supplement). After quality control procedures were performed to remove participants with missing data, and keeping only white British unrelated individuals, 121,604 individuals with AUDIT total scores were available. AUDIT total score was created by taking the sum of items 1–10, for all participants, including those who endorsed currently never drinking alcohol (as they could still endorse past hazardous use on items 9 and 10). We also created AUDIT subdomain scores by aggregating the scores from items 1–3, which include the information pertaining to alcohol consumption (AUDIT-C, $N=121,604$), and from items 4–10, which index the information pertaining to alcohol problems (AUDIT-P, $N=121,604$). These traits were \log_{10} -transformed to approximate a normal distribution (see Figure S2 in the online supplement).

Genotyping, Quality Control, and Imputation

Genotype imputation was performed on 487,409 individuals by the UK Biobank team, using IMPUTE4 (18) and the Haplotype Reference Consortium reference panel. After quality control, 16,213,998 single-nucleotide polymorphisms (SNPs) remained for GWAS analyses. Additional details on genotyping and quality control are provided in the Supplementary Methods section in the online supplement.

Discovery GWASs Using the UK Biobank Sample

GWAS analyses were performed using BGENIE, version 1.1 (18), with AUDIT scores (total score, AUDIT-C score, and AUDIT-P score, tested independently) as the outcome variable and age, sex, genotyping array, and the first 20 principal components derived from genotype data as covariates. (See the Supplementary Methods section in the online supplement for more details.) To identify independently associated variants (“index variants”), clump-based pruning was applied in FUMA (fuma.ctglab.nl) using an r^2 of 0.1 and a 1-Mb sliding window using the UK Biobank white British sample as the linkage disequilibrium (LD) reference panel. A 1-Mb window was used because of the regions of extended LD on chromosomes 4q23 and 17q21.31, which were associated with AUDIT scores in this study.

In addition, we performed a series of 18 case-control GWAS analyses of AUDIT total score using different thresholds (cases: $\geq 8, 10, 12, 15, 18, 20$; compared with controls: $\leq 2, 3, 4$). The sample size at each threshold is listed in Table S2 in the online supplement. The results of these analyses were used to determine which thresholds would produce the highest genetic correlation estimates with DSM-IV alcohol dependence (13).

SNP Heritability Analyses

The SNP heritability of UK Biobank AUDIT scores (total, AUDIT-C, AUDIT-P) was calculated using a genomic restricted maximum likelihood (GREML) method implemented in Genetic Complex Trait Analysis (GCTA) (19) on a subset of

117,072 unrelated individuals, using a relatedness cutoff of 0.05 and controlling for age and sex. GREML analyses were run using genotyped SNPs with a minor allele frequency greater than 0.01 to construct the genetic relationship matrix.

GWAS Meta-Analysis of AUDIT Total Score Using the UK Biobank and 23andMe Samples

Because the genetic correlation of AUDIT total score between the UK Biobank and 23andMe cohorts was high ($r_g=0.77$, $SE=0.12$, $p=7.15 \times 10^{-11}$), we performed a sample size–based meta-analysis of AUDIT total score from the UK Biobank and 23andMe cohorts using METAL (version 2011-03-25) (20). This meta-analysis comprises a total of 141,932 research participants of European ancestry and 9,519,872 genetic variants that passed quality control. We used clump-based pruning (see the “Discovery GWASs Using the UK Biobank” section, above) to identify independently associated variants. For each GWAS signal, we defined a set of credible variants using a Bayesian refinement method developed by Maller et al. (21). These credible sets are considered to have a 99% probability of containing the “causal” variant at each locus. Credible set analyses were performed in R (<https://github.com/hailianghuang/FM-summary>) for each of the index variants associated with AUDIT total score in the GWAS meta-analysis using SNPs within 1 Mb with an $r^2 > 0.4$ to the index variant. All downstream genetic analyses of AUDIT total score were performed using the GWAS meta-analysis summary statistics. The 23andMe AUDIT GWAS has been published previously (10), and 30,441 participants from the UK Biobank cohort were included in a previous GWAS of alcohol consumption (9). The GWAS of alcohol consumption in the UK Biobank sample involved individuals genotyped during the first UK Biobank data release (9). The individuals from the UK Biobank in the present study are those who completed the mental health questionnaire in 2017 and include those genotyped in both the first and second UK Biobank data releases; the sample overlap between these two data sets is 30,441 individuals.

Functional Mapping and Annotation of GWAS Meta-Analysis

We used FUMA, version 1.2.8 (22), to study the functional consequences of the index SNPs and of the SNPs contained in each credible set, which included categories from the ANNOVAR tool, combined annotation-dependent depletion (CADD) scores, RegulomeDB scores (www.regulomedb.org/), expression quantitative trait loci, and chromatin states. We also studied the regulatory consequences of the index SNPs using the Variant Effect Predictor tool (VEP; Ensembl GRCh37).

Gene-Set and Pathway Analyses

We performed MAGMA (22) competitive gene-set and pathway analyses using the summary statistics from the GWAS meta-analysis of AUDIT total score and the AUDIT-C and AUDIT-P score subsets. SNPs were mapped to 18,546 protein-coding genes

from Ensembl, build 85. Gene sets were obtained from Msigdb, version 5.2 (“Curated gene sets,” “GO terms”).

Gene-Based Association Using Transcriptomic Data With S-PrediXcan

We used S-PrediXcan (23) to predict gene expression levels in 10 brain tissues and to test whether the predicted gene expression correlates with AUDIT scores. We used precomputed tissue weights from the Genotype-Tissue Expression (version 7) project database (<https://www.gtexportal.org/>) as the reference transcriptome data set. Further details are provided in the Supplementary Methods section in the online supplement.

Genetic Correlation Analysis

We used LD score regression (LDSC) (github.com/bulik/ldsc; ldsc.broadinstitute.org) to identify genetic correlations between traits (24). This method was used to calculate genetic correlations (r_g) between AUDIT total score, AUDIT-C score, and AUDIT-P score, and 39 other traits and diseases (see Tables S3, S4, and S5 in the online supplement). We did not constrain the intercepts in our analysis, as we could not quantify the exact amount of sample overlap between cohorts. We used false discovery rate to correct for multiple testing (25). We also used LDSC to examine genetic correlations between various dichotomized versions of AUDIT and DSM-IV alcohol dependence (13). To test for significant differences between the genetic correlations, z-score statistics were calculated (see Table S6 in the online supplement).

RESULTS

UK Biobank Sample Characteristics

In the UK Biobank cohort, there were 121,604 individuals with AUDIT scores available for GWAS analysis (see Table S7 in the online supplement). The UK Biobank sample was 56.2% female ($N=68,389$), and the mean age was 56.1 years ($SD=7.7$). The mean AUDIT total score was 5.0 ($SD=4.18$, range=0–40); a histogram showing the distribution of the scores is provided in Figure S2 in the online supplement. Over the previous year, 91.9% of the participants reported drinking 1 or 2 drinks on a single drinking day. Over the previous year, 6.3% of the participants reported that they were not able to stop drinking once they started, and 10.7% felt guilt or remorse after drinking (see Table S7 in the online supplement). The mean AUDIT total score was significantly higher for males than for females (6.09 [$SD=4.45$] and 4.15 [$SD=3.72$], respectively; $\beta=0.47$, $p<2 \times 10^{-6}$) (see Figure S3 in the online supplement). In addition, age was negatively correlated with AUDIT scores ($\beta=-0.02$, $p<2 \times 10^{-6}$) (see Table S8 in the online supplement). Therefore, both sex and age were used as covariates in the GWAS analyses. The mean AUDIT-C score was 4.24 ($SD=2.83$) and the mean AUDIT-P score was 0.75 ($SD=2.0$). As expected, there was a moderate positive phenotypic correlation between AUDIT-C and AUDIT-P scores ($r=0.478$, 95% CI=0.473–0.481, $p<2 \times 10^{-16}$) (see Table S8 in the online supplement).

TABLE 1. Index SNPs Associated With AUDIT Scores in a GWAS Meta-Analysis of AUDIT Scores and Corresponding p Values in the UK Biobank and 23andMe Cohorts^a

Chromosome	Index Variant	Genes (Number of Credible SNPs)	UK Biobank (N=121,604)			
			A1	Beta	SE	p
2p21	rs4953148	<i>LINC01833</i> (17)	T	0.007	0.001	2.8×10^{-8}
2p23.3	rs1260326	<i>SNX17</i> (1), <i>GCKR</i> (3)	C	0.008	0.001	2.1×10^{-10}
3q25.33	rs1920650	<i>SMC4</i> , <i>TRIM59</i> , <i>B3GAT3P1</i> , <i>KPNA4</i> , <i>SCARNA7</i> , <i>KRT8P12</i> , <i>RPL6P8</i> , <i>ARL14</i> (132)	C	0.006	0.001	1.0×10^{-6}
4p14	rs11940694	<i>KLB</i> (7)	G	0.01	0.001	9.8×10^{-20}
4p14	rs4975012	<i>KLB</i> (6)	A	0.008	0.001	1.9×10^{-8}
4q23	rs146788033	<i>METAP1</i> (1)	G	-0.04	0.005	1.5×10^{-22}
4q23	rs11733695	<i>RP11-696N14.1</i> (1)	A	-0.07	0.006	9.1×10^{-26}
4q23	rs3114045	<i>ADH1C</i> (12)	C	0.01	0.002	3.4×10^{-10}
4q24	rs188514326	<i>RP11-588P8.1</i> (1)	C	-0.04	0.009	3.0×10^{-7}
4q24	rs13135092	<i>RN7SL728P</i> (1), <i>SLC39A8</i> (5)	G	-0.02	0.002	4.1×10^{-13}
8q22.1	rs35040843	<i>RP11-700E23.3</i> (30)	T	0.007	0.001	1.7×10^{-7}
10p11.23	rs7078436	<i>JCAD</i> (9)	G	-0.006	0.001	1.8×10^{-6}
11p11.2	rs2293576	<i>NDUFS3</i> (3), <i>SPI1</i> (6), <i>SLC39A13</i> (3), <i>PSMC3</i> (3), <i>RAPSN</i> (6), <i>CELF1</i> (16), <i>RP11-750H9.5</i> (15), <i>Y_RNA</i> (1)	A	0.007	0.001	5.0×10^{-9}
17q21.31	rs62062288	<i>CRHR1</i> (142), <i>DND1P1</i> (1), <i>ENSG00000262372</i> (1), <i>ENSG00000262500</i> (2), <i>ENSG00000262881</i> (12), <i>KANSL1</i> (376), <i>MAPT</i> (755), <i>MAPT-AS1</i> (255), <i>NSF</i> (15), <i>PLEKHM1</i> (1), <i>SPPL2C</i> (11), <i>STH</i> (1), <i>RP11-105N13.4</i> (541), <i>WNT3</i> (16)	A	-0.009	0.001	1.6×10^{-9}
19q13.33	rs492602	<i>FUT2</i> (29), <i>MAMSTR</i> (8), <i>RASIP1</i> (23), <i>IZUMO1</i> (6)	G	0.006	0.001	7.0×10^{-8}

^aProtein coding genes listed are those spanned by SNPs in each credible set. AUDIT=Alcohol Use Disorders Identification Test; GWAS=genome-wide association study; SNP=single-nucleotide polymorphism.

SNP Heritability in the UK Biobank Sample

We estimated the SNP heritability of AUDIT total score to be 12% (GCTA: SE=0.48%, $p=4.6 \times 10^{-273}$; LDSC: 8.6%, SE=0.50%), which is similar to the previously published AUDIT estimate (10). The SNP heritability for AUDIT-C score was 11% (GCTA: SE=0.47%, $p=1.5 \times 10^{-211}$; LDSC: 8.4%, SE=0.55%), and 9% for AUDIT-P score (GCTA: SE=0.46%, $p=2.0 \times 10^{-178}$; LDSC: 5.9%, SE=0.48%).

GWAS of AUDIT Scores in the UK Biobank Sample

The significant results ($p < 5 \times 10^{-8}$) of the GWAS of AUDIT total score in the UK Biobank cohort are presented in Table S9 in the online supplement; this analysis revealed 12 independent GWAS signals located in eight loci. The UK Biobank GWASs of AUDIT-C and AUDIT-P score subsets are summarized in Tables S10 and S11 and Figure S4 in the online supplement. Seven of these 12 independent GWAS signals were also significantly associated with AUDIT-C score; the same index variants were identified in the two analyses. An additional GWAS signal was also identified close to *FBNP4*. For AUDIT-P score, five independent GWAS signals were significantly associated, and these loci were also associated with the total AUDIT and AUDIT-C scores. The rs1229984 SNP in *ADH1B* was not available for meta-analysis in the 23andMe sample and was not in Hardy-Weinberg equilibrium in the UK Biobank sample used in the present study ($p=3.2 \times 10^{-16}$); however, in the total UK Biobank white British sample, there was no significant deviation from Hardy-Weinberg equilibrium ($p=0.13$). The associations

between rs1229984 and AUDIT scores are presented in Tables S9, S10, and S11 in the online supplement. The rs1229984 SNP was strongly associated with all AUDIT scores in the UK Biobank sample ($\beta=0.04$ – 0.06 , $p \leq 1.0 \times 10^{-45}$), but this SNP was not included for clump-based pruning and downstream analyses. We therefore performed a conditional analysis of the SNPs on 4q23 and 4q24 in the UK Biobank sample to determine whether any of these associations were significant after controlling for rs1229984 genotype. While rs13107325 on 4q24 remained significantly associated with AUDIT total score after controlling for rs1229984 genotype, the association between rs146788033, rs11733695, and rs3114045 and AUDIT total score became nonsignificant, suggesting that these loci are tagging the strong rs1229984 signal in this region.

GWAS Meta-Analysis of AUDIT Total Score

The GWAS meta-analysis of the UK Biobank and 23andMe samples found 15 independent GWAS signals (see Table S12 in the online supplement) associated with AUDIT total score spanning 10 genomic loci (Table 1). Figure 1 presents the Manhattan and quantile-quantile plots of the GWAS meta-analysis of AUDIT total score, and Figures S5–S14 in the online supplement present the regional association plots for the independent signals. The inflation factor of the meta-analysis GWAS was $\lambda_{GC}=1.22$ with an LDSC intercept of 1.008 (SE=0.007), suggesting that the majority of the inflation is due to polygenicity. The 15 independent SNPs show 100% sign concordance for association with AUDIT total

23andMe (N=20,328)				Meta-Analysis (N=141,932)			
A1	Beta	SE	p	A1	Z Score	p	Direction
T	0.03	0.036	3.5×10^{-1}	A	-5.59	2.3×10^{-8}	-/-
T	-0.10	0.034	2.7×10^{-3}	T	-7.15	8.7×10^{-13}	-/-
T	-0.07	0.037	4.2×10^{-2}	T	-5.63	1.7×10^{-8}	-/-
G	0.08	0.034	2.5×10^{-2}	A	-9.38	6.8×10^{-21}	-/-
G	-0.006	0.003	4.9×10^{-2}	A	5.94	2.8×10^{-9}	+/+
G	-0.03	0.008	1.4×10^{-4}	A	10.48	1.0×10^{-25}	+/+
G	0.05	0.012	6.7×10^{-6}	A	-11.42	3.4×10^{-30}	-/-
T	-0.15	0.048	1.6×10^{-3}	T	-7.18	7.2×10^{-13}	-/-
G	0.03	0.013	4.9×10^{-2}	C	-5.49	4.1×10^{-8}	-/-
G	-0.09	0.060	1.2×10^{-1}	A	7.59	3.2×10^{-14}	+/+
T	0.09	0.038	8.9×10^{-3}	T	5.55	2.9×10^{-8}	+/+
G	-0.10	0.035	3.1×10^{-3}	A	5.74	9.6×10^{-9}	+/+
G	-0.06	0.035	1.2×10^{-1}	A	5.83	5.5×10^{-9}	+/+
G	0.09	0.042	2.1×10^{-2}	A	-6.21	5.4×10^{-10}	-/-
G	0.05	0.033	1.7×10^{-1}	A	-5.57	2.5×10^{-8}	-/-

score across the UK Biobank and 23andMe samples (Table 1); 11 of these SNPs were nominally associated with AUDIT total score in the 23andMe sample ($p \leq 0.05$), and all index SNPs were associated with AUDIT total score in the UK Biobank sample ($p < 1.8 \times 10^{-6}$).

The top hit for the GWAS meta-analysis of AUDIT total score was a variant (rs11733695) located downstream (879 base pairs) from *ADH6* ($p = 3.4 \times 10^{-30}$), which is a member of the alcohol dehydrogenase gene family. The rs11733695 SNP is in low LD ($r^2 = 0.17$) with the functional SNP in *ADH1B*, rs1229984, which is known to alter alcohol metabolism (26). In addition, two other regions in 4q23 were associated with AUDIT total score in the meta-analysis: one of the index SNPs was located in the *ADH1C* gene; however, conditional analysis of this region in the UK Biobank sample alone suggests that these multiple hits may in fact be tagging the rs1229984 signal. This region has been previously associated with alcohol consumption, alcohol use disorders, and AUDIT scores (6, 7, 9, 10).


We also replicated the association between *KLB* (see Table S12 in the online supplement), on chromosome 4q14, and alcohol consumption (9, 11, 12); the index SNP rs11940694, which is located in the intron of *KLB*, was associated with AUDIT total score in the present study. Clump-based pruning identified rs11940694 and rs4975012 as independent hits in the *KLB* region. Credible set analysis suggests that rs2046330 is the index SNP in the region represented by rs4975012 (see Table S13 in the online supplement). AUDIT total score was also associated with SNPs that localized to *GCKR* on chromosome 2p23.3, which has been previously associated with alcohol consumption (9, 11). Five SNPs comprised the

credible set at the *GCKR* locus, which also spans the *SNX17* gene, including the missense variant (rs1260326) in *GCKR* that was identified as the index SNP.

We identified GWAS signals in several regions that have not been previously implicated in the genetics of alcohol use disorders, including 2p21, 17q21, 3q25, 8q22, 10p11, 11p11, and 19q13. The index SNP rs13135092 in the 4q24 region is located in an intron of *SLC39A8*; the remainder of the credible set for this locus is located in a noncoding pseudogene, *RN7SL728P*. *SLC39A8* is highly pleiotropic (27), but it is a novel association in relation to alcohol. A region of association on 2p21 contains 17 SNPs that are localized to the noncoding RNA *LINC01833*. A novel region of association was also detected on chromosome 10p11.23; this region contains nine credible SNPs that localize to the *JCAD* (junctional cadherin 5 associated) gene. *JCAD* encodes an endothelial cell junction protein and has previously been associated with coronary heart disease (28).

The remaining novel associations on 3q25, 8q, 11p11, 17q21, and 19q13.3 were more complex. The index variants on chromosome 8q22.1 were not localized to any genes, and it is unclear from the credible set analyses what the causal variants may be at these loci. The credible SNP sets for the 3q25.33, 11p11.2, 17q21.31, and 19q13.3 regions contained more than 50 SNPs each, which spanned several genes. For example, the index SNP on chromosome 17q21.31 was an intronic SNP in *MAPT*, which encodes the tau protein and has been robustly associated with Parkinson's disease (30, 31) (see Table S14 in the online supplement) and other neurodegenerative tauopathies (32) and, more recently, with

FIGURE 1. Manhattan and QQ Plots for the SNP-Based GWAS Meta-Analysis of AUDIT Total Score (N=141,932)^a



^a AUDIT=Alcohol Use Disorders Identification Test; GWAS=genome-wide association study; QQ=quantile-quantile; SNP=single-nucleotide polymorphism.

neuroticism (33). However, we note that the region of association on chromosome 17q21.31 spans the corticotropin receptor gene (*CRHR1*), which has been associated with alcohol use in animals and humans (34). Thus, because of the extended complex LD in this region, we are unable to determine the likely causal variant. Similarly, the index SNP (rs2293576) at chromosome 11p11.2 is a synonymous SNP of the zinc transporter gene *SLC39A13*; however, this region includes 54 associated SNPs, which map to five additional genes. Lastly, the index SNP on 19q13.3 is a synonymous variant in *FUT2*. *FUT2* encodes galactoside 2- α -L-fucosyltransferase 2, which controls the expression of ABO blood group antigens. However, 70 SNPs form this credible set, and they span four genes in total.

We used FUMA to functionally annotate all 1,290 SNPs in the credible sets (see Table S13 in the online supplement). The majority of the SNPs were intronic (76.9%; N=993) or intergenic (11.6%; N=149), and only 26 SNPs (2.0%) were exonic. Furthermore, 38 SNPs showed CADD scores >12.37, which is the suggested threshold to be considered deleterious (35). The exonic SNPs (rs601338, rs17651549, rs13107325) of *FUT2*, *MAPT*, and *SLC39A8*, respectively, had the highest CADD scores (>34), suggesting potential deleterious protein effects. Overall, 164 SNPs had RegulomeDB scores of 1a–1f, showing evidence of potential regulatory effects; 90.1% of the SNPs were in open chromatin regions (minimum chromatin state, 1–7).

Gene-Based and Pathway Analyses

We used MAGMA (22) to perform a gene-based association analysis, which identified 40 genes that were significantly associated with AUDIT total score ($p < 2.70 \times 10^{-6}$) (see Table S15 and Figure S15 in the online supplement). As expected, the majority of these genes were in the 10 GWAS loci (i.e., *KLB*, *GCKR*); *DRD2* and *CRHR1* were also among the top hits. In addition, the analysis revealed a strong burden signal in *CADM2* ($p = 1.64 \times 10^{-9}$), where the index variant in GWAS meta-analysis did not reach genome-wide

significance. We did not identify any canonical pathways that were significantly associated with AUDIT score (see Table S16 in the online supplement).


Gene-based (MAGMA) analyses for the AUDIT-C and AUDIT-P subsets (see Figures S16 and S17 in the online supplement) revealed evidence of overlap (see Figure S18 and Table S17 in the online supplement). Two genes (*KLB*, *CADM2*) were associated with all three AUDIT traits (AUDIT total score, AUDIT-C score, and AUDIT-P score). There was considerable overlap between AUDIT total score and AUDIT-C score, with 21 overlapping genes associated at the gene-based level. Only one gene, *DRD2*, was associated with both AUDIT total score and AUDIT-P score.

S-PrediXcan

S-PrediXcan identified a positive correlation ($p < 1.07 \times 10^{-6}$) between AUDIT total score and the predicted expression of 26 genes across multiple brain tissues (full results are presented in Table S18 in the online supplement), including *MAPT* (cerebellum), *FUT2* (caudate and nucleus accumbens), and *CRHR1-IT1* (putamen, cerebellum, hippocampus, anterior cingulate cortex). SNPs in the region of *MAPT* and *FUT2* were associated with AUDIT total score in the GWAS. *MAPT* (cerebellum), *FUT2* (nucleus accumbens), and *CRHR1-IT1* (caudate, cortex, nucleus accumbens, hypothalamus, hippocampus) were also associated with AUDIT-C score (see Table S19 in the online supplement). S-PrediXcan for AUDIT-C and AUDIT-P scores (see Table S20 in the online supplement) revealed that lower predicted *RFC1* expression in the cerebellum and hemisphere, respectively, was associated with higher scores on both the AUDIT-C ($p = 7.84 \times 10^{-7}$) and the AUDIT-P ($p = 1.54 \times 10^{-6}$).

Genetic Correlations

We used LDSC to evaluate evidence for genetic correlations between our three primary traits (AUDIT total score, AUDIT-C score, and AUDIT-P score) and numerous other traits for which GWAS summary statistics were available;

FIGURE 2. Genetic Correlations Between the Three AUDIT Phenotypes (Total Score, AUDIT-C Score, AUDIT-P Score) and Several Traits Measured in Independent Cohorts^a

^a More information on the various traits is provided in Tables S3–S5 in the online supplement. ADHD=attention deficit hyperactivity disorder; AUDIT=Alcohol Use Disorders Identification Test; AUDIT-C=AUDIT items 1–3 (consumption); AUDIT-P=AUDIT items 4–10 (problematic consequences of drinking); BMI=body mass index.

^b AUDIT-P compared with AUDIT-C, $p < 0.01$, false discovery rate 5%.

^c AUDIT-P compared with AUDIT-C, $p < 0.05$.

* $p < 0.05$. ** $p < 0.01$. *** $p < 0.0001$.


these included alcohol and substance use traits, personality and behavioral traits, psychiatric disorders, blood lipid levels, and brain structure volumes (see Figure 2; see also Tables S3–S5 in the online supplement).

As expected, AUDIT-C and AUDIT-P scores were highly genetically correlated ($r_g = 0.70$, $p = 4.1 \times 10^{-70}$). AUDIT scores (AUDIT total score, AUDIT-C score, and AUDIT-P score) showed strong genetic correlations with alcohol consumption from two other studies ($r_g = 0.76$ – 0.96 , $p < 2.3 \times 10^{-9}$). Many of the genetic correlations with AUDIT-P score were significantly different from the correlations with AUDIT-C score (see Table S6 in the online supplement). AUDIT-C score had a significantly stronger ($p = 8.02 \times 10^{-3}$) correlation with alcohol consumption ($r_g = 0.92$, $p = 7.0 \times 10^{-164}$) than did

AUDIT-P score ($r_g = 0.76$, $p = 2.7 \times 10^{-52}$). In contrast, AUDIT total score and AUDIT-C score were only modestly correlated with alcohol dependence ($r_g = 0.39$ and 0.33 , respectively, $p < 8.2 \times 10^{-5}$), whereas AUDIT-P score showed a nominally stronger genetic correlation with alcohol dependence ($r_g = 0.63$, $p = 1.8 \times 10^{-8}$; AUDIT-P compared with AUDIT-C, $p = 0.033$) (see Table S6).

We detected positive genetic correlations between AUDIT scores (AUDIT total, AUDIT-C score, AUDIT-P score) and other substance use phenotypes, including lifetime smoking ($r_g = 0.24$ – 0.41 , $p < 1.6 \times 10^{-5}$) and cannabis use ($r_g = 0.26$ – 0.46 , $p < 1.1 \times 10^{-4}$). We also observed a positive genetic correlation between AUDIT-P score and cigarettes per day ($r_g = 0.28$, $p = 4.0 \times 10^{-3}$).

FIGURE 3. Dichotomizing AUDIT Total Score to Approximate DSM-IV Alcohol Dependence^a



^a Continuous AUDIT total score was used to categorize participants as case and control subjects at different thresholds; GWAS was performed on each, and the genetic correlation with DSM-IV alcohol dependence was calculated. Thresholds for AUDIT-determined cases were 8 (N=25,423), 10 (N=15,151), 12 (N=9,130), 15 (N=2,099), and 20 (N=1,290). The orange line is a visualization of the number of cases used at each threshold, corresponding to the N on the right-hand y-axis. AUDIT=Alcohol Use Disorders Identification Test; GWAS=genome-wide association study.

Several psychiatric disorders and related traits were positively genetically correlated with AUDIT-P score, including schizophrenia ($r_g=0.22$, $p=3.0 \times 10^{-10}$), bipolar disorder ($r_g=0.26$, $p=1.5 \times 10^{-4}$), ADHD ($r_g=0.23$, $p=1.1 \times 10^{-5}$), and major depressive disorder ($r_g=0.26$, $p=50.6 \times 10^{-3}$). Intriguingly, AUDIT-C score was negatively correlated with major depressive disorder ($r_g=-0.23$, $p=3.7 \times 10^{-3}$) and ADHD ($r_g=-0.10$, $p=1.8 \times 10^{-2}$), whereas AUDIT-P score showed positive genetic correlations with these same disease traits ($r_g=0.26$, $p=5.6 \times 10^{-3}$; $r_g=0.24$, $p=1.1 \times 10^{-5}$). We observed a positive genetic correlation between AUDIT-P score and depressive symptoms ($r_g=0.30$, $p=3.0 \times 10^{-8}$) and neuroticism ($r_g=0.18$, $p=2.6 \times 10^{-4}$) and a negative genetic correlation with subjective well-being ($r_g=-0.24$, $p=4.0 \times 10^{-5}$).

We observed positive genetic correlations between AUDIT total score and AUDIT-C score and education, college completion, and cognitive ability ($r_g=0.19-0.24$, $p<1.5 \times 10^{-5}$). The genetic correlations between AUDIT-P and education and college completion were near zero and were significantly lower than AUDIT-C score and AUDIT total score or education traits (see Table S6 in the online supplement).

There were negative genetic correlations between AUDIT total score and AUDIT-C score and obesity ($r_g=-0.16-0.17$, $p<1.1 \times 10^{-5}$), similar to previous reports regarding AUDIT total score (10) and alcohol consumption (9). In contrast, obesity was not significantly genetically correlated with AUDIT-P score ($r_g=0.01$, $p=0.90$). Similarly, HDL cholesterol and triglyceride levels were genetically correlated with AUDIT total score and AUDIT-C score ($r_g=0.19-0.22$, $p<9.3 \times 10^{-5}$, $r_g=-0.16$, $p<1.0 \times 10^{-4}$, respectively), but this association was not found for AUDIT-P score ($r_g=0.11$,

$p=2.2 \times 10^{-2}$, $r_g=-0.03$, $p=6.4 \times 10^{-1}$). Obesity showed significantly different correlations with both AUDIT-P and AUDIT-C scores (see Table S6).

Dichotomizing AUDIT Total Score to More Closely Approximate DSM-IV Alcohol Dependence

As AUDIT can be rapidly ascertained in large populations, we explored methods for dichotomizing AUDIT total score to optimize the genetic correlation with DSM-IV alcohol dependence (13). Higher genetic correlations with alcohol dependence were observed as the control threshold was increased from 2 to 4, and with increasingly stringent case cutoffs (Figure 3; see also Table S2 in the online supplement). The highest genetic correlation was observed for cases with AUDIT total score ≥ 20 and controls ≤ 4 ($r_g=0.90$, $SE=0.25$, $p=3.0 \times 10^{-4}$); however, this highly stringent threshold produced very few cases (N=1,290). The standard error of the estimate is much larger at more stringent case thresholds, and therefore these estimates should be interpreted with caution. Defining cases as ≥ 12 yielded an r_g of 0.82 ($SE=0.18$, $p=3.2 \times 10^{-6}$) while retaining more than seven times as many cases (N=9,130); these genetic correlations were not significantly different from those obtained using cases ≥ 20 and controls ≤ 4 ($p=0.80$).

DISCUSSION

We have presented the largest GWAS meta-analysis of AUDIT total score to date, using large population-based cohorts from the UK Biobank and 23andMe. We identified novel associations with AUDIT total score; the genes located

in these regions include *JCAD* and *SLC39A8*. We found evidence for association in several loci previously associated with alcohol use via single-variant and gene-based analyses (i.e., *KLB*, *GCKR*, *CADM2*). The SNP heritability of all AUDIT phenotypes ranged from 9% to 12%, demonstrating that common genetic factors account for a prominent proportion of the variation in alcohol use phenotypes. Furthermore, we showed that there is shared genetic architecture between AUDIT scores and other alcohol and substance use phenotypes. AUDIT-P score showed a positive genetic correlation with several psychiatric diseases, distinguishing AUDIT-P score from AUDIT-C score. Finally, using LDSC, we identified thresholds for dichotomizing AUDIT total score (score ≥ 12 to define cases and ≤ 4 to define controls) that maximize the genetic correlation with alcohol dependence while retaining a large number of participants.

Our top GWAS hits replicated previous association signals for alcohol use traits. The strongest associations with AUDIT score in this study spanned the alcohol metabolism genes on chromosome 4q23 (10). Variants in this region were associated with AUDIT total score, AUDIT-C score, and AUDIT-P score, demonstrating that alcohol metabolism is a risk factor for both alcohol consumption and problematic use. The second strongest signal, also associated with the three AUDIT phenotypes, is located in *KLB*, confirming the robust association of this gene with alcohol consumption in both humans (9, 11, 12) and mice (12). However, the biology of this locus could be more complex than previously described. Although the credible set analysis suggested that the more probable causal variants are all located on the first intron of *KLB*, one of these variants, rs11940694, is an expression quantitative trait locus for *RFC1* expression in the brain, and *S-PrediXcan* analysis predicted that lower expression of *RFC1* in the brain is associated with higher predicted AUDIT (AUDIT-C and AUDIT-P) scores. Interestingly, a gene in the complex GWAS signal on chromosome 19, Fibroblast growth factor 21 (*FGF21*), was associated with AUDIT scores (AUDIT total score, AUDIT-C score, AUDIT-P score) at the gene-based level (see Table S17 in the online supplement). *FGF21* regulates sweet and alcohol preference in mice as part of a receptor complex with β -Klotho (*KLB*) in the central nervous system (33). Additionally, we replicated the association between rs1260326 in the gene *GCKR* and alcohol consumption (9, 11), here associated with AUDIT total score and AUDIT-C score. Other loci previously associated with alcohol consumption include *CADM2* (9), which was associated at the gene-based level for all three AUDIT traits. Here, the burden analysis suggests that multiple (rare and common) variants are necessary to explain the association signal. Intriguingly, several of the novel associations with AUDIT scores were mapped to highly pleiotropic genes (*MAPT*, *FUT2*, *SLC39A8*) (27).

Genetic analysis of the AUDIT subsets revealed evidence of distinct genetic architecture between AUDIT-C and AUDIT-P (alcohol consumption versus problem use), with support from the gene-based (see Figures S16 and S17 in the

online supplement), *S-PrediXcan* (see Tables S19 and S20 in the online supplement), and genetic correlation analyses (see Figure 2). Furthermore, AUDIT-P score showed a strong genetic correlation with alcohol dependence (13). In contrast, AUDIT-C score had a stronger genetic correlation with alcohol consumption. Thus, partitioning AUDIT scores into different subsets (alcohol consumption versus problem use) may disentangle genetic factors that contribute to different aspects of vulnerability to alcohol use disorders.

Genetic overlap was observed for all measures of AUDIT and other substance use traits, including lifetime tobacco and cannabis use, as we previously reported (10, 36, 37), demonstrating that genetic risk factors for high AUDIT scores overlap with increased consumption of multiple drug types.

We found several significant differences between the genetic correlations with AUDIT-P and AUDIT-C scores. These differences were particularly pronounced for psychiatric and behavioral traits. AUDIT-P score was positively genetically correlated with psychopathology (schizophrenia, bipolar disorder, major depressive disorder, ADHD), personality traits (including neuroticism), and regional brain volumes. These associations have previously been observed at the phenotypic level; alcohol use disorders commonly co-occur in individuals with schizophrenia (38), bipolar disorder (39), major depressive disorder (40), and adult ADHD (41). Intriguingly, genetic risk for high AUDIT-C score was *negatively* correlated with major depressive disorder and ADHD, demonstrating that a distinct genetic component of AUDIT-P is shared with genetic risk for psychiatric disease. Regional volume abnormalities in subcortical brain regions of individuals with alcohol use disorders have been reported (42–44), although it is unclear whether these alterations are a result of high alcohol consumption or a preexisting susceptibility. We identified a positive genetic correlation between AUDIT-P score and increased caudate volume; however, the majority of studies report reductions in regional brain volumes associated with alcohol use disorders.

For AUDIT total score and AUDIT-C score, we showed positive genetic correlations with educational attainment and cognitive ability and negative genetic correlations with obesity, consistent with earlier reports (9, 10). These associations were not observed for AUDIT-P score. Similarly, HDL cholesterol levels showed a significant positive correlation, and triglyceride levels a negative correlation, with AUDIT total score and AUDIT-C score, but not AUDIT-P score. These patterns were previously observed for alcohol consumption (9). We could speculate that these differences may be linked to socioeconomic status. Alcohol consumption is often higher in individuals with higher socioeconomic status (45), whereas alcohol-related problems, such as binge drinking (46) and alcohol-related mortality (47), are more prevalent in individuals with lower socioeconomic status. Furthermore, individuals with low socioeconomic status are more likely to have alcohol use disorders with psychiatric comorbidities (48). Consistent with this idea, we find positive genetic correlations between AUDIT-C score and

education, a trait correlated with socioeconomic status (49), and positive genetic correlations between AUDIT-P score and psychopathology. Our findings provide further evidence that different dimensions of alcohol use associate differently with behavior and that these differences may have a biological underpinning.

A clinical diagnosis of an alcohol use disorder is often required to define cases for genetic studies. An alternative strategy would be to use AUDIT to infer alcohol use disorder case status; however, it has not been clear whether and how to perform meta-analyses between AUDIT scores and alcohol dependence. A GWAS meta-analysis for AUDIT score and alcohol dependence could be simplified if a threshold could be used to define cases and controls based on AUDIT score, an approach that was used by Mbarek et al. (50). We have provided empirical evidence about genetic correlations between AUDIT score and alcohol dependence using dichotomized AUDIT scores and found thresholds for AUDIT score that produced high genetic correlations with alcohol use disorders (see Figure 3). Genetic correlations increased as the upper threshold for cases was made more stringent, although the standard errors for all of these estimates were overlapping. The genetic correlation with alcohol dependence appeared to become asymptotic when case status was defined as ≥ 12 ; therefore, this threshold could be used to define case status. We also considered various thresholds for defining controls and found that a threshold of ≤ 4 produced a high genetic correlation with alcohol dependence while also retaining the largest number of subjects.

Our study is not without limitations. The AUDIT specifically asks about the past year, and thus it may not capture information on lifetime alcohol use and misuse. This is suboptimal for genetic studies, because it effectively measures a recent state rather than a stable trait. Measures capturing drinking and alcohol use disorders across the lifespan may be preferable. Also, although the mean AUDIT-C score was 4.24, the mean AUDIT-P score was considerably lower (0.75). Thus, we were not able to perform a more refined categorization (e.g., three subsets: consumption [items 1–3], dependence [items 4–6], hazardous use [items 7–10]), as fewer individuals endorsed the items comprising the AUDIT-P (see Table S7 in the online supplement). Furthermore, our study uses data from the UK Biobank and 23andMe research participants, who were volunteers not ascertained for alcohol use disorders, and hence our findings may not generalize to populations that show higher rates of alcohol use and dependence. Additional alcohol-related phenotypes (e.g., age at first use and patterns of alcohol drinking, including binge drinking) could be used in subsequent genetic studies to identify additional sources of genetic vulnerability for alcohol use disorders. Lastly, we offered guidelines to identify cases to use in genetic studies of alcohol use disorders (i.e., an AUDIT score ≥ 12) based on genetic correlations; however, these recommendations are not intended to determine thresholds for diagnosing dependence in a clinical setting. Future studies will be able to

test whether using the AUDIT as a surrogate for alcohol use disorders will be beneficial for gene discovery. In addition, several studies have argued that lower thresholds should be used for females, which has not been addressed in the present study.

CONCLUSIONS

We have reported the largest GWAS of AUDIT ever undertaken. We replicated previously identified signals (i.e., *ADH1B*, *ADH1C*; *KLB*; *GCKR*) and identified novel GWAS signals (i.e., *JACD*, *SLC39A8*) associated with AUDIT scores. We showed that different portions of the AUDIT (AUDIT-C, AUDIT-P) correlate with distinct traits, which will aid in dissecting genetic vulnerability to alcohol use and dependence. The genetic factors that predispose to high alcohol consumption inevitably overlap with those for problem drinking, as heavy drinking is generally a prerequisite for the development of hazardous use and dependence. However, not everyone who consumes alcohol experiences the same level of harmful consequences. By studying the different subsets of AUDIT, we identified genetic factors that may be specific to problem drinking. Larger studies of cohorts with a wider range of AUDIT-P scores are required to both replicate and expand these findings. Finally, we described an alternative strategy to rigorous ascertainment for genetic studies of alcohol use disorders, namely, an AUDIT score ≥ 12 to define cases and a score ≤ 4 to define controls, which could be used to achieve large sample sizes in a cost-efficient manner.

AUTHOR AND ARTICLE INFORMATION

From the Department of Psychiatry and the Institute for Genomic Medicine, University of California San Diego, La Jolla; 23andMe, Inc., Mountain View, Calif.; the Division of Psychiatry, the Department of Psychology, and the Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, U.K.; the Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis; and the Translational Research Laboratories, Center for Neurobiology and Behavior, Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia.

Address correspondence to Dr. Clarke (toni.clarke@ed.ac.uk).

The collaborators for the 23andMe Research Team are follows: Michelle Agee, Babak Alipanahi, Adam Auton, Robert K. Bell, Katarzyna Bryc, Sarah L. Elson, Pierre Fontanillas, Nicholas A. Furlotte, David A. Hinds, Karen E. Huber, Aaron Kleinman, Nadia K. Litterman, Jennifer C. McCreight, Matthew H. McIntyre, Joanna L. Mountain, Elizabeth S. Noblin, Carrie A.M. Northover, Steven J. Pitts, J. Fah Sathirapongsasuti, Olga V. Sazonova, Janie F. Shelton, Suyash Shringarpure, Chao Tian, Joyce Y. Tung, Vladimir Vacic, and Catherine H. Wilson.

Data availability: Full GWAS summary statistics for the 23andMe data set will be made available through 23andMe to qualified researchers under an agreement with 23andMe that protects the privacy of the 23andMe participants. Interested investigators should e-mail dataset-request@23andme.com for more information. GWAS summary statistics for the UK Biobank GWAS of AUDIT scores will be available on request.

Dr. Sanchez-Roige was supported by the Frontiers of Innovation Scholars Program (3-P3029), the Interdisciplinary Research Fellowship in Neuro-AIDS (MH081482), and a pilot award from DA037844. This research was conducted using the UK Biobank Resource, application number 4844, and

was supported by a Wellcome Trust Strategic Award (Stratifying Resilience and Depression Longitudinally; reference 104036/Z/14/Z) and by the Centre for Cognitive Ageing and Cognitive Epidemiology, which was funded by the Medical Research Council and the Biotechnology and Biological Sciences Research Council (reference MR/K026992/1).

The authors thank the research participants and employees of 23andMe for making this work possible.

Drs. Fontanillas and Elson and the members of the 23andMe Research Team are employees of 23andMe. Dr. McIntosh has received research support from Eli Lilly, Janssen, Pfizer, and the Sackler Foundation. The other authors report no financial relationships with commercial interests.

Received April 3, 2018; revision received July 18, 2018; accepted August 9, 2018; published online Oct. 19, 2018.

REFERENCES

- Heath AC, Madden PAF, Bucholz KK, et al: Genetic differences in alcohol sensitivity and the inheritance of alcoholism risk. *Psychol Med* 1999; 29:1069–1081
- Verhulst B, Neale MC, Kendler KS: The heritability of alcohol use disorders: a meta-analysis of twin and adoption studies. *Psychol Med* 2015; 45:1061–1072
- Frank J, Cichon S, Treutlein J, et al: Genome-wide significant association between alcohol dependence and a variant in the ADH gene cluster. *Addict Biol* 2012; 17:171–180
- Kilcoyne B, Shmulewitz D, Meyers JL, et al: Alcohol consumption mediates the relationship between ADH1B and DSM-IV alcohol use disorder and criteria. *J Stud Alcohol Drugs* 2014; 75:635–642
- Quillen EE, Chen X-D, Almasy L, et al: ALDH2 is associated to alcohol dependence and is the major genetic determinant of “daily maximum drinks” in a GWAS study of an isolated rural Chinese sample. *Am J Med Genet B Neuropsychiatr Genet* 2014; 165B: 103–110
- Gelernter J, Kranzler HR, Sherva R, et al: Genome-wide association study of alcohol dependence: significant findings in African- and European-Americans including novel risk loci. *Mol Psychiatry* 2014; 19:41–49
- Bierut LJ, Goate AM, Breslau N, et al: ADH1B is associated with alcohol dependence and alcohol consumption in populations of European and African ancestry. *Mol Psychiatry* 2012; 17:445–450
- Macgregor S, Lind PA, Bucholz KK, et al: Associations of ADH and ALDH2 gene variation with self report alcohol reactions, consumption, and dependence: an integrated analysis. *Hum Mol Genet* 2009; 18:580–593
- Clarke T-K, Adams MJ, Davies G, et al: Genome-wide association study of alcohol consumption and genetic overlap with other health-related traits in UK Biobank (N=112 117). *Mol Psychiatry* 2017; 22: 1376–1384
- Sanchez-Roige S, Fontanillas P, Elson SL, et al: Genome-wide association study of Alcohol Use Disorder Identification Test (AUDIT) scores in 20 328 research participants of European ancestry. *Addict Biol* (Epub ahead of print, Oct 23, 2017)
- Jorgenson E, Thai KK, Hoffmann TJ, et al: Genetic contributors to variation in alcohol consumption vary by race/ethnicity in a large multi-ethnic genome-wide association study. *Mol Psychiatry* 2017; 22:1359–1367
- Schumann G, Liu C, O'Reilly P, et al: KLB is associated with alcohol drinking, and its gene product β -Klotho is necessary for FGF21 regulation of alcohol preference. *Proc Natl Acad Sci USA* 2016; 113: 14372–14377
- Walters RK, Adams MJ, Adkins AE, et al: Trans-ancestral GWAS of alcohol dependence reveals common genetic underpinnings with psychiatric disorders. *bioRxiv* 257311; March 10, 2018 (doi: <https://doi.org/10.1101/257311>)
- Saunders JB, Babor TF: AUDIT questionnaire. *Addiction* 1993; 88:1–2
- Babor TF: The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Care, 2nd ed. Geneva, World Health Organization, 2001
- Bulik-Sullivan BK, Loh P-R, Finucane HK, et al: LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 2015; 47:291–295
- Allen N, Sudlow C, Downey P, et al: UK Biobank: current status and what it means for epidemiology. *Health Policy Technol* 2012; 1:123–126
- Bycroft C, Freeman C, Petkova D, et al: Genome-wide genetic data on ~500,000 UK Biobank participants. *bioRxiv* 166298; July 20, 2017 (<https://www.biorxiv.org/content/early/2017/07/20/166298>)
- Yang J, Lee SH, Goddard ME, et al: GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet* 2011; 88:76–82
- Willer CJ, Li Y, Abecasis GR: METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010; 26:2190–2191
- Maller JB, McVean G, Byrnes J, et al: Bayesian refinement of association signals for 14 loci in 3 common diseases. *Nat Genet* 2012; 44:1294–1301
- Watanabe K, Taskesen E, van Bochoven A, et al: Functional mapping and annotation of genetic associations with FUMA. *Nat Commun* 2017; 8:1826
- Barbeira A, Shah KP, Torres JM, et al: Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics. *bioRxiv* 045260; March 23, 2016 (<https://www.biorxiv.org/content/early/2017/10/03/045260>)
- Bulik-Sullivan B, Finucane HK, Anttila V, et al: An atlas of genetic correlations across human diseases and traits. *Nat Genet* 2015; 47: 1236–1241
- Benjamini Y, Hochberg MT: Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* 1995; 57:289–300
- Hurley TD, Edenberg HJ: Genes encoding enzymes involved in ethanol metabolism. *Alcohol Res* 2012; 34:339–344
- Pickrell JK, Berisa T, Liu JZ, et al: Detection and interpretation of shared genetic influences on 42 human traits. *Nat Genet* 2016; 48:709–717
- Erdmann J, Willenborg C, Nahrstaedt J, et al: Genome-wide association study identifies a new locus for coronary artery disease on chromosome 10p11.23. *Eur Heart J* 2011; 32:158–168
- Bustamante M, Standl M, Bassat Q, et al: A genome-wide association meta-analysis of diarrhoeal disease in young children identifies FUT2 locus and provides plausible biological pathways. *Hum Mol Genet* 2016; 25:4127–4142
- Satake W, Nakabayashi Y, Mizuta I, et al: Genome-wide association study identifies common variants at four loci as genetic risk factors for Parkinson's disease. *Nat Genet* 2009; 41:1303–1307
- Simón-Sánchez J, Schulte C, Bras JM, et al: Genome-wide association study reveals genetic risk underlying Parkinson's disease. *Nat Genet* 2009; 41:1308–1312
- Pittman AM, Fung HC, de Silva R: Untangling the tau gene association with neurodegenerative disorders. *Hum Mol Genet* 2006; 15:R188–R195
- Talukdar S, Owen BM, Song P, et al: FGF21 regulates sweet and alcohol preference. *Cell Metab* 2016; 23:344–349
- Zorrilla EP, Logrip ML, Koob GF: Corticotropin releasing factor: a key role in the neurobiology of addiction. *Front Neuroendocrinol* 2014; 35:234–244
- Kircher M, Witten DM, Jain P, et al: A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet* 2014; 46:310–315
- Nivard MG, Verweij KJH, Minică CC, et al: Connecting the dots: genome-wide association studies in substance use. *Mol Psychiatry* 2016; 21:733–735
- Vink JM, Hottenga JJ, de Geus EJC, et al: Polygenic risk scores for smoking: predictors for alcohol and cannabis use? *Addiction* 2014; 109:1141–1151
- Drake RE, Mueser KT: Co-occurring alcohol use disorder and schizophrenia. *Alcohol Res Health* 2002; 26:99–102
- Sonne SC, Brady KT: Substance abuse and bipolar comorbidity. *Psychiatr Clin North Am* 1999; 22:609–627, ix

40. Petrakis IL, Gonzalez G, Rosenheck R, et al: Comorbidity of alcoholism and psychiatric disorders: an overview. *Alcohol Res Health* 2002; 26:81–89
41. Ohlmeier MD, Peters K, Te Wildt BT, et al: Comorbidity of alcohol and substance dependence with attention-deficit/hyperactivity disorder (ADHD). *Alcohol Alcohol* 2008; 43:300–304
42. Pfefferbaum A, Sullivan EV, Mathalon DH, et al: Longitudinal changes in magnetic resonance imaging brain volumes in abstinent and relapsed alcoholics. *Alcohol Clin Exp Res* 1995; 19:1177–1191
43. Pfefferbaum A, Lim KO, Zipursky RB, et al: Brain gray and white matter volume loss accelerates with aging in chronic alcoholics: a quantitative MRI study. *Alcohol Clin Exp Res* 1992; 16:1078–1089
44. Cardenas VA, Studholme C, Meyerhoff DJ, et al: Chronic active heavy drinking and family history of problem drinking modulate regional brain tissue volumes. *Psychiatry Res* 2005; 138:115–130
45. Collins SE: Associations between socioeconomic factors and alcohol outcomes. *Alcohol Res* 2016; 38:83–94
46. Fone DL, Farewell DM, White J, et al: Socioeconomic patterning of excess alcohol consumption and binge drinking: a cross-sectional study of multilevel associations with neighbourhood deprivation. *BMJ Open* 2013; 3:3
47. Probst C, Roerecke M, Behrendt S, et al: Socioeconomic differences in alcohol-attributable mortality compared with all-cause mortality: a systematic review and meta-analysis. *Int J Epidemiol* 2014; 43:1314–1327
48. Lee SJ, Sudore RL, Williams BA, et al: Functional limitations, socioeconomic status, and all-cause mortality in moderate alcohol drinkers. *J Am Geriatr Soc* 2009; 57:955–962
49. Marioni RE, Davies G, Hayward C, et al: Molecular genetic contributions to socioeconomic status and intelligence. *Intelligence* 2014; 44:26–32
50. Mbarek H, Milaneschi Y, Fedko IO, et al: The genetics of alcohol dependence: twin and SNP-based heritability, and genome-wide association study based on AUDIT scores. *Am J Med Genet B Neuropsychiatr Genet* 2015; 168:739–748