

TABLE S1. Self-reported medication information for MDD-medicated participants

MDD-medicated participant	Primary antidepressant medication	Primary medication dose	Time on primary medication (weeks)	Other reported medication
1	venlafaxine	75 mg	227	lamotrigine 150 mg
2	venlafaxine	75 mg	16	trazodone 200mg
3	duloxetine	60 mg	328	--
4	citalopram	40 mg	143	--
5	venlafaxine	150 mg	205	lamotrigine 200 mg bupropion 300 mg
6	--	--	--	--
7	bupropion	--	308	--
8	bupropion	300 mg	119	lithium 300 mg
9	citalopram	--	76	--
10	--	--	--	--
11	escitalopram	10 mg	27	bupropion 150 mg
12	sertraline	--	1.5	--
13	bupropion	300 mg	10	--
14	escitalopram	10 mg	--	--
15	citalopram	20 mg	126	--
16	duloxetine	--	7	--
17	venlafaxine	300 mg	106	--
18	escitalopram	10 mg	81	--
19	citalopram	20 mg	24	--
20	duloxetine	60 mg	13	--
21	paroxetine	20 mg	520	--

FIGURE S1. An example of a single study trial with dysphoric (upper left), neutral (upper right), positive (lower right), and threat (lower left) stimuli. To avoid copyright infringement, IAPS images used in the study have been replaced with similar public domain images in this figure.



FIGURE S2. Mean total gaze duration for positive stimuli, by 5-second epoch and group

