


# New Spring Titles

## from American Psychiatric Association Publishing



### The American Psychiatric Publishing Textbook of Suicide Risk Assessment and Management, Third Edition

Edited by Liza H. Gold, M.D., and Richard L. Frierson, M.D.

2020 • 477 pages • ISBN 978-1-61537-223-2  
Hardcover • \$120.00 • Item #37223

2020 • 477 pages • ISBN 978-1-61537-284-3  
eBook • \$96.00 • Item #37284

### Clinical Manual of Youth Addictive Disorders

Edited by Yifrah Kaminer, M.D., M.B.A., and Ken C. Winters, Ph.D.

2020 • 632 pages • ISBN 978-1-61537-236-2  
Paperback • \$75.00 • Item #37236

2020 • 632 pages • ISBN 978-1-61537-281-2  
eBook • \$60.00 • Item #37281

### Psychotherapeutic Approaches to Sexual Problems

An Essential Guide for Mental Health Professionals

Stephen B. Levine, M.D.

2020 • 157 pages • ISBN 978-1-61537-283-6  
Paperback • \$39.00 • Item #37283

2020 • 157 pages • ISBN 978-1-61537-285-0  
eBook • \$31.00 • Item #37285

### Practical Strategies in Geriatric Mental Health

Cases and Approaches

Edited by Laura B. Dunn, M.D., and Erin L. Cassidy-Eagle, Ph.D.

2020 • 512 pages • ISBN 978-1-61537-148-8  
Paperback • \$75.00 • Item #37148

2020 • 512 pages • ISBN 978-1-61537-257-7  
eBook • \$60.00 • Item #37257

### Clinical Manual of Pediatric Consultation-Liaison Psychiatry, Second Edition

Richard J. Shaw, M.B., B.S., and David R. DeMaso, M.D.

2020 • 566 pages • ISBN 978-1-61537-231-7  
Paperback • \$75.00 • Item #37231

2020 • 566 pages • ISBN 978-1-61537-279-9  
eBook • \$60.00 • Item #37279

### Divorce and Co-parenting

A Support Guide for the Modern Family

Elissa P. Benedek, M.D., and Samantha A. Huettner, J.D.

2020 • 272 pages • ISBN 978-1-61537-220-1  
Paperback • \$29.00 • Item #37220

2020 • 272 pages • ISBN 978-1-61537-271-3  
eBook • \$23.00 • Item #37271

### Behavioral and Psychological Symptoms of Dementia

Art Walaszek, M.D.

2020 • 284 pages • ISBN 978-1-61537-168-6  
Paperback • \$65.00 • Item #37168

2020 • 284 pages • ISBN 978-1-61537-267-6  
eBook • \$52.00 • Item #37267

### Obsessive-Compulsive Personality Disorder

Edited by Jon E. Grant, M.D., M.P.H., J.D., Samuel R. Chamberlain, MBBChir, Ph.D., MRCPsych, and Anthony Pinto, Ph.D.

2020 • 275 pages • ISBN 978-1-61537-224-9  
Paperback • \$54.00 • Item #37224

2020 • 275 pages • ISBN 978-1-61537-280-5  
eBook • \$43.00 • Item #37280

### Combating Physician Burnout

A Guide for Psychiatrists

Edited by Sheila LoboPrabhu, M.D., Richard F. Summers, M.D., and H. Steven Moffic, M.D.

2020 • 344 pages • ISBN 978-1-61537-227-0  
Paperback • \$60.00 • Item #37227

2020 • 344 pages • ISBN 978-1-61537-272-0  
eBook • \$48.00 • Item #37272

### Learning Supportive Psychotherapy

An Illustrated Guide, Second Edition  
Core Competencies in Psychotherapy

Arnold Winston, M.D., Richard N. Rosenthal, M.D., and Laura Weiss Roberts, M.D., M.A.

2020 • 232 pages • ISBN 978-1-61537-234-8  
Paperback • \$70.00 • Item #37234

2020 • 232 pages • ISBN 978-1-61537-287-4  
eBook • \$56.00 • Item #37287

### Clinical Manual of Supportive Psychotherapy, Second Edition

Peter N. Novalis, M.D., Ph.D., Virginia Singer, D.N.P., and Roger Peele, M.D.

2020 • 469 pages • ISBN 978-1-61537-165-5  
Paperback • \$77.00 • Item #37165

2020 • 469 pages • ISBN 978-1-61537-273-7  
eBook • \$62.00 • Item #37273

### Physician Well-Being

Cases and Solutions

Peter Yellowlees, MBBS, M.D.

2020 • 265 pages • ISBN 978-1-61537-240-9  
Paperback • \$55.00 • Item #37240

2020 • 265 pages • ISBN 978-1-61537-290-4  
eBook • \$44.00 • Item #37290

AMERICAN  
PSYCHIATRIC  
ASSOCIATION  
PUBLISHING



ORDER @ [www.appi.org](http://www.appi.org)

Email: [appi@psych.org](mailto:appi@psych.org) | Toll Free: 1-899-368-5777



20% Discount for American Psychiatric Association Members

25% Discount for APA Resident-Fellow Members

## Subscription and Business Information

*The American Journal of Psychiatry*, ISSN 0002-953X, is published monthly by the American Psychiatric Association.

The American Psychiatric Association does not hold itself responsible for statements made in its publications by contributors or advertisers. Unless so stated, material in *The American Journal of Psychiatry* does not reflect the endorsement, official attitude, or position of the American Psychiatric Association or of the Journal's Editorial Board.

Periodicals postage paid at Washington, DC, and additional mailing offices. POSTMASTER: Send address changes to *The American Journal of Psychiatry*, Circulation Department, American Psychiatric Association, 800 Maine Avenue, S.W., Suite 900, Washington, D.C. 20024.

### Editorial Office

*American Journal of Psychiatry*  
American Psychiatric Association  
800 Maine Avenue, S.W., Suite 900  
Washington, D.C. 20024  
E-mail: [ajp@psych.org](mailto:ajp@psych.org)

### Customer Service

Phone: (800) 368-5777 (toll-free)  
E-mail: [appi@psych.org](mailto:appi@psych.org)

### APA Member Subscription Services

Phone: (888) 35-PSYCH (toll-free)

### Subscriptions

Regular and resident-fellow members in the U.S. receive print and online access as a member benefit; subscriptions for non-members start at \$348 per year. Online only subscriptions start at \$315 per year. Go to <http://www.appi.org/products> for more information.

Institutional subscriptions are tier priced. For institutional site license or pricing information, contact (202) 559-3729 or [institutions@psych.org](mailto:institutions@psych.org).

*Additional subscription options, including single issues and student rates*

Contact Customer Service

### Advertising

*Pharmaceutical Print and Online Advertising*  
Frank Cox, Kathleen Harrison, Tim Wolfinger  
Pharmaceutical Media, Inc.  
30 East 33rd Street  
New York, NY 10016  
Phone: (212) 904-0379  
E-mail: [twolfinger@pminy.com](mailto:twolfinger@pminy.com)

*Nonpharmaceutical and Online Sales*  
Eamon Wood  
Pharmaceutical Media, Inc.  
E-mail: [ewood@pminy.com](mailto:ewood@pminy.com)  
(see address and phone above).

### Permissions and Reprints

Material published in the journals of the American Psychiatric Association and American Psychiatric Association Publishing is protected by copyright and all rights are reserved. Material may not be reproduced in any form or by any means without written permission from the copyright owner. For permission to reproduce material published by APA, please visit <http://www.appi.org/customer-service/permissions> for more information. Permission can also be secured through the Copyright Clearance Center ([www.copyright.com](http://www.copyright.com)).

APA does not require that permission be obtained for the photocopying of isolated articles for nonprofit classroom or library reserve use; all fees associated with such permission are waived. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale. Requests for commercial distribution and bulk reprints should be directed to (202) 609-7075.

Indexed in Abstracts for Social Workers, Academic Abstracts, Biological Abstracts, Chemical Abstracts, Chicago Psychoanalytic Literature Index, Cumulative Index to Nursing Literature, Excerpta Medica, Hospital Literature Index, Index Medicus, International Nursing Index, Nutrition Abstracts, Psychological Abstracts, Science Citation Index, Social Science Source, and Social Sciences Index.

Pages are produced by Sheridan Journal Services (Waterbury, VT) and printed by Sheridan NH (Hanover, NH) on acid-free paper effective with Volume 169, Number 1, January 2012.

© 2020 American Psychiatric Association.

# INGREZZA® (valbenazine) capsules

for oral use

**Brief Summary:** for full Prescribing Information and Patient Information, refer to package insert.

## INDICATION AND USAGE

INGREZZA® (valbenazine) capsules is indicated for the treatment of adults with tardive dyskinesia.

## CONTRAINDICATIONS

INGREZZA is contraindicated in patients with a history of hypersensitivity to valbenazine or any components of INGREZZA. Rash, urticaria, and reactions consistent with angioedema (e.g., swelling of the face, lips, and mouth) have been reported.

## WARNINGS AND PRECAUTIONS

### Somnolence

INGREZZA can cause somnolence. Patients should not perform activities requiring mental alertness such as operating a motor vehicle or operating hazardous machinery until they know how they will be affected by INGREZZA.

### QT Prolongation

INGREZZA may prolong the QT interval, although the degree of QT prolongation is not clinically significant at concentrations expected with recommended dosing. In patients taking a strong CYP2D6 or CYP3A4 inhibitor, or who are CYP2D6 poor metabolizers, INGREZZA concentrations may be higher and QT prolongation clinically significant. For patients who are CYP2D6 poor metabolizers or are taking a strong CYP2D6 inhibitor, dose reduction may be necessary. For patients taking a strong CYP3A4 inhibitor, reduce the dose of INGREZZA to 40 mg once daily. INGREZZA should be avoided in patients with congenital long QT syndrome or with arrhythmias associated with a prolonged QT interval. For patients at increased risk of a prolonged QT interval, assess the QT interval before increasing the dosage.

### Parkinsonism

INGREZZA may cause parkinsonism in patients with tardive dyskinesia. Parkinsonism has also been observed with other VMAT2 inhibitors. In the 3 placebo-controlled clinical studies in patients with tardive dyskinesia, the incidence of parkinson-like adverse events was 3% of patients treated with INGREZZA and <1% of placebo-treated patients. Postmarketing safety reports have described parkinson-like symptoms, some of which were severe and required hospitalization. In most cases, severe parkinsonism occurred within the first 2 weeks after starting or increasing the dose of INGREZZA. Associated symptoms have included falls, gait disturbances, tremor, drooling, and hypokinesia. In cases in which follow-up clinical information was available, parkinson-like symptoms were reported to resolve following discontinuation of INGREZZA therapy. Reduce the dose or discontinue INGREZZA treatment in patients who develop clinically significant parkinson-like signs or symptoms.

## ADVERSE REACTIONS

The following adverse reactions are discussed in more detail in other sections of the labeling:

- Hypersensitivity
- Somnolence
- QT Prolongation
- Parkinsonism

### Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### Variable and Fixed Dose Placebo-Controlled Trial Experience

The safety of INGREZZA was evaluated in 3 placebo-controlled studies, each 6 weeks in duration (fixed dose, dose escalation, dose reduction), including 445 patients. Patients were 26 to 84 years of age with moderate to severe tardive dyskinesia and had concurrent diagnoses of mood disorder (27%) or schizophrenia/schizoaffective disorder (72%). The mean age was 56 years. Patients were 57% Caucasian, 39% African-American, and 4% other. With respect to ethnicity, 28% were Hispanic or Latino. All subjects continued previous stable regimens of antipsychotics; 85% and 27% of subjects, respectively, were taking atypical and typical antipsychotic medications at study entry.

#### Adverse Reactions Leading to Discontinuation of Treatment

A total of 3% of INGREZZA treated patients and 2% of placebo-treated patients discontinued because of adverse reactions.

#### Common Adverse Reactions

Adverse reactions that occurred in the 3 placebo-controlled studies at an incidence of ≥2% and greater than placebo are presented in Table 1.

**Table 1: Adverse Reactions in 3 Placebo-Controlled Studies of 6-week Treatment Duration Reported at ≥2% and >Placebo**

Adverse Reaction <sup>1</sup>	INGREZZA (n=262) (%)	Placebo (n=183) (%)
<b>General Disorders</b>		
Somnolence (somnolence, fatigue, sedation)	10.9%	4.2%
<b>Nervous System Disorders</b>		
Anticholinergic effects (dry mouth, constipation, disturbance in attention, vision blurred, urinary retention)	5.4%	4.9%
Balance disorders/fall (fall, gait disturbance, dizziness, balance disorder)	4.1%	2.2%
Headache	3.4%	2.7%
Akathisia (akathisia, restlessness)	2.7%	0.5%
<b>Gastrointestinal Disorders</b>		
Vomiting	2.6%	0.6%
Nausea	2.3%	2.1%
<b>Musculoskeletal Disorders</b>		
Arthralgia	2.3%	0.5%

<sup>1</sup> Within each adverse reaction category, the observed adverse reactions are listed in order of decreasing frequency.

### Other Adverse Reactions Observed During the Premarketing Evaluation of INGREZZA

Other adverse reactions of ≥1% incidence and greater than placebo are shown below. The following list does not include adverse reactions: 1) already listed in previous tables or elsewhere in the labeling, 2) for which a drug cause was remote, 3) which were so general as to be uninformative, 4) which were not considered to have clinically significant implications, or 5) which occurred at a rate equal to or less than placebo.

**Endocrine Disorders:** blood glucose increased

**General Disorders:** weight increased

**Infectious Disorders:** respiratory infections

**Neurologic Disorders:** drooling, dyskinesia, extrapyramidal symptoms (non-akathisia)

**Psychiatric Disorders:** anxiety, insomnia

During controlled trials, there was a dose-related increase in prolactin. Additionally, there was a dose-related increase in alkaline phosphatase and bilirubin, suggesting a potential risk for cholestasis.

### Postmarketing Experience

The following adverse reactions have been identified during post-approval use of INGREZZA that are not included in other sections of labeling. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Immune System Disorders:** hypersensitivity reactions (including allergic dermatitis, angioedema, pruritis, and urticaria)

**Skin and Subcutaneous Tissue Disorders:** rash

## DRUG INTERACTIONS

### Drugs Having Clinically Important Interactions with INGREZZA

**Table 2: Clinically Significant Drug Interactions with INGREZZA**

<b>Monoamine Oxidase Inhibitors (MAOIs)</b>	
<b>Clinical Implication:</b>	Concomitant use of INGREZZA with MAOIs may increase the concentration of monoamine neurotransmitters in synapses, potentially leading to increased risk of adverse reactions such as serotonin syndrome, or attenuated treatment effect of INGREZZA.
<b>Prevention or Management:</b>	Avoid concomitant use of INGREZZA with MAOIs.
<b>Examples:</b>	isocarboxazid, phenelzine, selegiline
<b>Strong CYP3A4 Inhibitors</b>	
<b>Clinical Implication:</b>	Concomitant use of INGREZZA with strong CYP3A4 inhibitors increased the exposure (C <sub>max</sub> and AUC) to valbenazine and its active metabolite compared with the use of INGREZZA alone. Increased exposure of valbenazine and its active metabolite may increase the risk of exposure-related adverse reactions.
<b>Prevention or Management:</b>	Reduce INGREZZA dose when INGREZZA is coadministered with a strong CYP3A4 inhibitor.
<b>Examples:</b>	itraconazole, ketoconazole, clarithromycin
<b>Strong CYP2D6 Inhibitors</b>	
<b>Clinical Implication:</b>	Concomitant use of INGREZZA with strong CYP2D6 inhibitors may increase the exposure (C <sub>max</sub> and AUC) to valbenazine's active metabolite compared with the use of INGREZZA alone. Increased exposure of active metabolite may increase the risk of exposure-related adverse reactions.
<b>Prevention or Management:</b>	Consider reducing INGREZZA dose based on tolerability when INGREZZA is coadministered with a strong CYP2D6 inhibitor.
<b>Examples:</b>	paroxetine, fluoxetine, quinidine
<b>Strong CYP3A4 Inducers</b>	
<b>Clinical Implication:</b>	Concomitant use of INGREZZA with a strong CYP3A4 inducer decreased the exposure of valbenazine and its active metabolite compared to the use of INGREZZA alone. Reduced exposure of valbenazine and its active metabolite may reduce efficacy.
<b>Prevention or Management:</b>	Concomitant use of strong CYP3A4 inducers with INGREZZA is not recommended.
<b>Examples:</b>	rifampin, carbamazepine, phenytoin, St. John's wort <sup>1</sup>
<b>Digoxin</b>	
<b>Clinical Implication:</b>	Concomitant use of INGREZZA with digoxin increased digoxin levels because of inhibition of intestinal P-glycoprotein (P-gp).
<b>Prevention or Management:</b>	Digoxin concentrations should be monitored when coadministering INGREZZA with digoxin. Increased digoxin exposure may increase the risk of exposure-related adverse reactions. Dosage adjustment of digoxin may be necessary.

<sup>1</sup> The induction potency of St. John's wort may vary widely based on preparation.

### Drugs Having No Clinically Important Interactions with INGREZZA

Dosage adjustment for INGREZZA is not necessary when used in combination with substrates of CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2E1, or CYP3A4/5 based on *in vitro* study results.

## OVERDOSAGE

### Human Experience

The pre-marketing clinical trials involving INGREZZA in approximately 850 subjects do not provide information regarding symptoms with overdose.

### Management of Overdosage

No specific antidotes for INGREZZA are known. In managing overdose, provide supportive care, including close medical supervision and monitoring, and consider the possibility of multiple drug involvement. If an overdose occurs, consult a Certified Poison Control Center (1-800-222-1222 or www.poison.org).

For further information on INGREZZA, call 84-INGREZZA (844-647-3992).

Distributed by:  
Neurocrine Biosciences, Inc.  
San Diego, CA 92130

INGREZZA is a registered trademark of Neurocrine Biosciences, Inc.  
CP-VBZ-US-0203v4 07/19



IN ADULT PATIENTS WITH TARDIVE DYSKINESIA (TD)

## Choose INGREZZA for results you can see<sup>1</sup>

INGREZZA<sup>®</sup> (valbenazine) capsules reduced TD severity at 6 weeks,  
with results you can start to see as early as 2 weeks<sup>1-3</sup>



See how INGREZZA reduced TD severity in patients at [INGREZZAHCP.com/results](https://www.INGREZZAHCP.com/results)

### Important Information

#### INDICATION & USAGE

INGREZZA<sup>®</sup> (valbenazine) capsules is indicated for the treatment of adults with tardive dyskinesia.

#### IMPORTANT SAFETY INFORMATION

##### CONTRAINDICATIONS

INGREZZA is contraindicated in patients with a history of hypersensitivity to valbenazine or any components of INGREZZA. Rash, urticaria, and reactions consistent with angioedema (e.g., swelling of the face, lips, and mouth) have been reported.

##### WARNINGS & PRECAUTIONS

###### Somnolence

INGREZZA can cause somnolence. Patients should not perform activities requiring mental alertness such as operating a motor vehicle or operating hazardous machinery until they know how they will be affected by INGREZZA.

###### QT Prolongation

INGREZZA may prolong the QT interval, although the degree of QT prolongation is not clinically significant at concentrations expected with recommended dosing. INGREZZA should be avoided in patients with congenital long QT syndrome or with arrhythmias associated with a prolonged QT interval. For patients at increased risk of a prolonged QT interval, assess the QT interval before increasing the dosage.

##### WARNINGS & PRECAUTIONS (continued)

###### Parkinsonism

INGREZZA may cause parkinsonism in patients with tardive dyskinesia. Parkinsonism has also been observed with other VMAT2 inhibitors. Reduce the dose or discontinue INGREZZA treatment in patients who develop clinically significant parkinson-like signs or symptoms.

##### ADVERSE REACTIONS

The most common adverse reaction ( $\geq 5\%$  and twice the rate of placebo) is somnolence. Other adverse reactions ( $\geq 2\%$  and  $>$ Placebo) include: anticholinergic effects, balance disorders/falls, headache, akathisia, vomiting, nausea, and arthralgia.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch at [www.fda.gov/medwatch](https://www.fda.gov/medwatch) or call 1-800-FDA-1088.

**Please see the adjacent page for Brief Summary of Prescribing Information and visit [www.INGREZZAHCP.com/PI](https://www.INGREZZAHCP.com/PI) for full Prescribing Information.**