

Supplemental Table A. Pooled Means on the Adult Self-Rated Level 1 Cross-Cutting Measures

Level 1 item	Range of Scores	Mean Score (SD)
Depression 1	0-4	1.98 (0.38)
Depression 2	0-4	2.09 (0.30)
Anger	0-4	1.86 (0.34)
Mania 1	0-4	1.29 (0.41)
Mania 2	0-4	0.83 (0.35)
Anxiety 1	0-4	2.21 (0.35)
Anxiety 2	0-4	1.46 (0.34)
Anxiety 3	0-4	1.85 (0.37)
Somatic Distress 1	0-4	1.61 (0.38)
Somatic Distress 2	0-4	1.25 (0.36)
Suicide	0-4	0.65 (0.25)
Psychosis 1	0-4	0.55 (0.23)
Psychosis 2	0-4	0.41 (0.23)
Sleep	0-4	2.08 (0.35)
Memory	0-4	1.54 (0.35)
Repetitive Thoughts	0-4	1.51 (0.37)
Repetitive Behaviors	0-4	0.98 (0.34)
Dissociation	0-4	1.05 (0.33)
Personality 1	0-4	1.51 (0.38)
Personality 2	0-4	1.86 (0.37)
Substance Use 1—Alcohol	0-4	0.34 (0.21)
Substance Use 2—Tobacco	0-4	1.15 (0.21)
Substance Use 3—Other Substances	0-4	0.35 (0.23)

Supplemental Table B. Pooled Means on the Parent- and Child-Rated Level 1 Cross-Cutting Measures

Level 1 Item	Range of Scores	Mean Score (SD)		
		Parent of Child < 11 years	Parent of Child 11+ years	Child 11+ years
Depression 1	0-4	1.04 (0.35)	1.19 (0.34)	1.20 (0.34)
Depression 2	0-4	0.93 (0.27)	1.33 (0.31)	1.10 (0.31)
Irritability	0-4	1.77 (0.36)	1.79 (0.32)	1.48 (0.37)
Anger	0-4	2.18 (0.32)	1.80 (0.31)	1.28 (0.34)
Mania 1	0-4	0.72 (0.31)	0.62 (0.28)	0.67 (0.33) ^b
Mania 2	0-4	0.89 (0.34)	0.63 (0.30)	1.06 (0.44)
Anxiety 1	0-4	1.51 (0.32)	1.33 (0.37)	1.28 (0.32)
Anxiety 2	0-4	1.18 (0.35)	1.22 (0.36)	1.09 (0.34)
Anxiety 3	0-4	0.85 (0.35)	0.80 (0.33)	0.76 (0.34)
Somatic Distress 1	0-4	1.11 (0.26)	1.44 (0.30)	1.28 (0.30)
Somatic Distress 2	0-4	0.65 (0.25)	0.85 (0.27)	0.78 (0.30)
Psychosis 1	0-4	0.16 (0.13)	0.18 (0.14)	0.32 (0.23) ^b
Psychosis 2	0-4	0.14 (0.14)	0.16 (0.16)	0.30 (0.25)
Sleep	0-4	1.76 (0.38)	1.71 (0.31)	1.52 (0.40)
Attention	0-4	2.52 (0.37)	2.20 (0.34)	1.66 (0.40) ^b
Repetitive Thoughts 1	0-4	0.51 (0.28)	0.35 (0.22)	0.59 (0.32)
Repetitive Behaviors 1	0-4	0.38 (0.20)	0.40 (0.22)	0.76 (0.31)

Repetitive Thoughts 2	0-4	0.42 (0.23)	0.37 (0.21)	0.46 (0.20)
Repetitive Behaviors 2	0-4	0.31 (0.19)	0.27 (0.23)	0.55 (0.28)
Substance Use 1—Alcohol	0-2 (P) 0-1 (C)	a	0.11 (0.10)	—
Substance Use 2—Tobacco	0-2 (P) 0-1 (C)	a	0.14 (0.07)	—
Substance Use 3—Illegal Drugs	0-2 (P) 0-1 (C)	a	0.09 (0.10)	—
Substance Use 4—Legal Drugs	0-2 (P) 0-1 (C)	0.01 (0.07) ^a	0.04 (0.07)	—
Suicide 1	0-2 (P) 0-1 (C)	0.20 (0.16)	0.18 (0.12)	—
Suicide 2	0-2 (P) 0-1 (C)	0.12 (0.07) ^b	0.25 (0.12)	—

^a Summary statistics could not be computed for these items at the following sites because all responses were identical within the site:

Baystate: Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 3

Colorado: Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 3

Columbia: Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 4

Stanford: Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 4, Suicide 2

^b Summary statistics were not included in the pooled estimates for these items at the following sites because standard errors for the ICC estimates were greater than 0.1:

Columbia: Parent respondents (child <11): Substance Use 3

Stanford: Parent respondents (child <11): Substance Use 3

Baystate: Child respondents Mania 1

Columbia: Child respondents Psychosis 1 and Attention

- Mean score and standard deviation are not provided since responses are Yes/No

Supplemental Table C. Pooled Means on the Adult Self-Rated Combined Level 1 and Level 2 Cross-Cutting Measures

Cross-Cutting Domain	Range of Scores	Mean Score (SD)
Depression	0-6	3.27 (0.43)
Anger	0-6	2.56 (0.52)
Mania	0-6	1.43 (0.44)
Anxiety	0-6	3.21 (0.42)
Somatic Distress	0-6	2.35 (0.47)
Sleep	0-6	3.12 (0.52)
Repetitive Thoughts and Behaviors	0-6	1.26 (0.33)
Substance Use—Other Substances	0-5	0.29 (0.26)

Supplemental Table D. Pooled Means on the Parent- and Child-Rated Combined Level 1 and Level 2 Cross-Cutting Measures

Cross-Cutting Domain	Range of Scores	Mean Score (SD)		
		Parent of Child <11 years	Parent of Child 11+ years	Child 11+ years
Depression	0-6	1.70 (0.43)	2.02 (0.45)	1.96 (0.43)
Anger	0-6	3.04 (0.46)	2.68 (0.43)	2.34 (0.52)
Irritability	0-6 (P)	2.50 (0.30)	2.17 (0.30)	1.73 (0.35)
	0-4 (C)			
Mania	0-6	1.34 (0.41)	1.04 (0.37)	1.45 (0.50)
Anxiety	0-6	2.15 (0.41)	1.92 (0.42)	1.87 (0.44)
Somatic Symptoms	0-6	1.42 (0.32)	1.85 (0.36)	1.85 (0.42)
Sleep	0-6	2.29 (0.49)	2.37 (0.43)	2.08 (0.57)
Inattentiveness	0-5 (P)	3.31 (0.46)	2.89 (0.45)	—

Repetitive Thoughts and Behaviors	0-6 (C)	–	–	1.48 (0.44)
Substance Use 1—Alcohol	0-5	a	0.04 (0.05) ^b	0.16 (0.12)
Substance Use 2—Tobacco	0-5	a	0.15 (0.13) ^a	0.26 (0.11)
Substance Use 3—Illegal Drug Use	0-5	a,b	0.03 (0.06) ^{a,b}	0.12 (0.09)
Substance Use 4—Legal Drug Use	0-5	0.01 (0.03) ^a	0.01 (0.03) ^{a,b}	0.05 (0.10) ^a
Suicidal Ideation	0-2 (P)	0.11 (0.11)	0.11 (0.09) ^a	0.27 (0.29)
	0-8 (C)			
Suicide Attempt	0-3	0.08 (0.07) ^a	0.15 (0.08)	0.29 (0.10)

–indicates that the item was not assessed in that patient group

^a Summary statistics could not be computed for these items at the following sites because all response were identical within the site:

Baystate: Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 3

Colorado: Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 3

Columbia: Child respondents: Substance Use 4

Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 4

Parent respondents (child 11+): Substance Use 2, Substance Use 4, Suicide Ideation

Stanford: Parent respondents (child <11): Substance Use 1, Substance Use 2, Substance Use 4, Suicide Attempt

Parent respondents (child 11+): Substance Use 3

^b Summary statistics were not included in the pooled estimates for these items at the following sites because standard errors for the ICC estimates were greater than 0.1:

Baystate: Parent respondents (child 11+): Substance Use 3, Substance Use 4

Columbia: Parent respondents (child <11): Substance Use 3

Parent respondents (child 11+): Substance Use 1, Substance Use 3

Stanford: Parent respondents (child <11): Substance Use 3

Supplemental Table E. Pooled Means on the Clinician-Rated Cross-Cutting Measures for Adults and Children

Cross-Cutting Domain	Range of Scores	Mean Score (SD)	
		Adult Patient	Child Patient
		18+ years	6-17 years
Suicide	0-4	0.72 (0.27)	0.42 (0.25)
Psychosis	0-4	0.56 (0.25)	0.06 (0.10) ^a

^a Summary statistics were not included in the pooled estimates for the psychosis rating at Stanford, Columbia and Colorado because standard errors for the ICC estimates were greater than 0.1.