

Table 1: Prevalence of Odd Thinking in Borderline Patients and Axis II Comparison Subject over 16 Years of Prospective Follow-up												
	BL	2 YR FU	4 YR FU	6 YR FU	8 YR FU	10 YR FU	12 YR FU	14 YR FU	16 YR FU	RRR Diagnosis Time	95%CI Diagnosis Time	P-value Diagnosis Time
Any Odd Thinking												
BPD	85.5 (248)	65.8 (181)	48.0 (129)	33.0 (87)	25.9 (66)	26.1 (65)	35.3 (86)	32.8 (78)	37.2 (86)	1.92 0.26	1.55, 2.39 0.22, 0.32	<0.001 <0.001
OPD	54.2 (39)	29.9 (20)	20.3 (13)	14.3 (9)	8.1 (5)	8.2 (5)	11.7 (7)	15.3 (9)	15.5 (9)			
Marked Superstitiousness												
BPD	21.4 (62)	13.8 (38)	11.9 (32)	14.4 (38)	6.3 (16)	2.8 (7)	9.4 (23)	13.0 (31)	12.12 (28)	1.96 0.50	1.11, 3.47 0.36, 0.70	NS <0.001
OPD	12.5 (9)	6.0 (4)	3.1 (2)	6.4 (4)	4.8 (3)	1.6 (1)	1.7 (1)	5.1 (3)	3.5 (2)			
Magical Thinking												
BPD	16.6 (48)	10.9 (30)	10.0 (27)	7.6 (20)	7.5 (19)	7.2 (18)	6.2 (15)	7.6 (18)	8.2 (19)	4.13 0.43	1.74, 9.79 0.26, 0.69	0.001 <0.001
OPD	6.9 (5)	1.5 (1)	1.6 (1)	1.6 (1)	0.0 (0)	0.0 (0)	0.0 (0)	1.7 (1)	0.0 (0)			
Sixth Sense												
BPD	21.7 (63)	13.1 (36)	15.2 (41)	13.3 (35)	5.9 (15)	6.8 (17)	7.0 (17)	8.4 (20)	8.7 (20)	2.06 0.33	1.16, 3.82 0.22, 0.49	NS <0.001
OPD	11.1 (8)	4.5 (3)	6.3 (4)	6.4 (4)	3.2 (2)	1.6 (1)	3.3 (2)	1.7 (1)	3.6 (2)			
Telepathy												
BPD	4.5 (13)	1.5 (4)	4.1 (11)	2.7 (7)	0.8 (2)	1.6 (4)	1.2 (3)	0.0 (0)	1.3 (3)	1.55 0.18	0.38, 6.30 0.07, 0.48	NS 0.001
OPD	2.8 (2)	1.5 (1)	1.6 (1)	1.6 (1)	1.6 (1)	0.0 (0)	0.0 (0)	0.0 (0)	1.7 (2)			
Clairvoyance												
BPD	12.8 (37)	7.3 (20)	3.7 (10)	4.6 (12)	1.2 (3)	1.6 (4)	3.7 (9)	2.1 (5)	3.9 (9)	1.37 0.20	0.60, 3.13 0.10, 0.39	NS <0.001
OPD	4.2 (3)	6.0 (4)	4.7 (3)	4.8 (3)	3.2 (2)	0.0 (0)	1.7 (1)	3.4 (2)	1.7 (1)			
Overvalued Ideas												
BPD	75.2 (218)	54.2 (149)	30.9 (83)	19.3 (51)	14.5 (37)	18.9 (47)	24.6 (60)	21.0 (50)	21.2 (49)	2.10 0.16	1.64, 2.68 0.12, 0.21	<0.001 <0.001
OPD	43.1 (31)	20.9 (14)	12.5 (8)	4.8 (3)	3.2 (2)	8.2 (5)	8.3 (5)	6.8 (4)	6.9 (4)			

Table 2: Prevalence of Unusual Perceptual Experiences in Borderline Patients and Axis II Comparison Subject over 16 Years of Prospective Follow-up

	BL	2 YR FU	4 YR FU	6 YR FU	8 YR FU	10 YR FU	12 YR FU	14 YR FU	16 YR FU	RRR Diagnosis Time	95%CI Diagnosis Time	P-value Diagnosis Time
Any Unusual Perceptions												
BPD	76.2 (221)	58.2 (160)	47.2 (127)	34.1 (90)	32.2 (82)	27.3 (68)	25.8 (63)	24.0 (57)	26.4 (61)	2.27 0.24	1.66, 3.12 0.20, 0.30	<0.001 <0.001
OPD	27.8 (20)	31.3 (21)	12.5 (8)	7.9 (5)	12.9 (8)	9.8 (6)	6.7 (4)	1.7 (1)	1.7 (1)			
Recurrent Illusions												
BPD	32.8 (95)	17.5 (48)	10.8 (29)	11.7 (31)	7.5 (19)	4.8 (12)	8.2 (20)	6.3 (15)	5.2 (12)	3.55 0.13	1.99, 6.35 0.84, 0.21	<0.001 <0.001
OPD	4.2 (3)	7.5 (5)	3.1 (2)	4.8 (3)	3.2 (2)	4.9 (3)	0.0 (0)	0.0 (0)	0.0 (0)			
Depersonalization												
BPD	59.7 (173)	48.7 (134)	39.0 (105)	26.1 (69)	28.6 (73)	24.1 (60)	18.4 (45)	18.5 (44)	22.1 (51)	2.27 0.26	1.57, 3.28 0.21, 0.33	<0.001 <0.001
OPD	25.0 (18)	20.9 (14)	10.9 (7)	4.8 (3)	9.7 (6)	4.9 (3)	3.3 (2)	1.7 (1)	1.7 (1)			
Derealization												
BPD	60.3 (175)	45.1 (124)	36.4 (98)	25.8 (68)	22.8 (58)	17.7 (44)	15.6 (38)	17.7 (42)	20.4 (47)	3.49 0.22	2.24, 5.44 0.17, 0.29	<0.001 <0.001
OPD	16.7 (12)	14.9 (10)	4.7 (3)	4.8 (3)	4.8 (3)	4.9 (3)	5.0 (3)	0.0 (0)	1.7 (1)			

Table 3: Prevalence of Paranoid Experiences in Borderline Patients and Axis II Comparison Subject over 16 Years of Prospective Follow-up

	BL	2 YR FU	4 YR FU	6 YR FU	8 YR FU	10 YR FU	12 YR FU	14 YR FU	16 YR FU	RRR Diagnosis Time	95%CI Diagnosis Time	P-value Diagnosis Time
Any Nondelusional Paranoia												
BPD	86.6 (251)	69.5 (191)	62.1 (167)	57.6 (152)	45.1 (115)	42.6 (106)	43.9 (107)	38.2 (91)	43.3 (100)	2.01 0.42	1.62, 2.51 0.37, 0.48	<0.001 <0.001
OPD	43.1 (31)	38.8 (26)	34.4 (22)	22.2 (14)	24.2 (15)	21.3 (13)	13.3 (8)	18.6 (11)	20.7 (12)			
Undue Suspiciousness												
BPD	71.4 (207)	58.6 (161)	54.3 (146)	46.2 (122)	35.3 (90)	31.7 (79)	35.7 (87)	31.9 (76)	37.2 (86)	2.29 0.43	1.74, 3.02 0.37, 0.50	<0.001 <0.001
OPD	29.2 (21)	26.9 (18)	28.1 (18)	17.5 (11)	21.0 (13)	13.1 (8)	11.7 (7)	17.0 (10)	19.0 (11)			
Ideas of Reference												
BPD	61.7 (179)	43.3 (119)	37.9 (102)	33.0 (87)	27.5 (70)	24.5 (61)	24.6 (60)	21.0 (50)	28.6 (66)	2.28 0.35	1.69, 3.06 0.29, 0.43	<0.001 <0.001
OPD	27.8 (20)	20.9 (14)	10.9 (7)	14.3 (9)	11.3 (7)	13.1 (8)	6.7 (4)	8.5 (5)	6.9 (4)			
Other Paranoid Ideation												
BPD	51.7 (150)	34.9 (96)	33.8 (91)	23.5 (62)	12.2 (31)	13.3 (33)	13.5 (33)	16.4 (39)	17.8 (41)	3.06 0.24	2.25, 4.58 0.19, 0.33	<0.001 <0.001
OPD	9.7 (7)	11.9 (8)	14.1 (9)	9.5 (6)	12.9 (8)	9.8 (6)	5.0 (3)	5.1 (3)	10.3 (6)			

Table 4: Prevalence of Quasi-Psychotic Thought in Borderline Patients and Axis II Comparison Subject over 16 Years of Prospective Follow-up												
	BL	2 YR FU	4 YR FU	6 YR FU	8 YR FU	10 YR FU	12 YR FU	14 YR FU	16 YR FU	RRR Diagnosis Time	95%CI Diagnosis Time	P-value Diagnosis Time
Any Quasi-psychotic Thought												
BPD	56.6 (164)	34.9 (96)	33.8 (91)	19.7 (52)	8.6 (22)	13.3 (33)	13.9 (34)	5.9 (14)	7.4 (17)	3.24 0.09	2.00, 5.24 0.07, 0.13	<0.001 <0.001
OPD	19.4 (14)	9.0 (6)	6.3 (4)	6.4 (4)	3.2 (2)	3.3 (2)	5.0 (3)	1.7 (1)	0.0 (0)			
Quasi Delusions												
BPD	48.6 (141)	25.8 (71)	25.7 (69)	14.8 (39)	5.5 (14)	3.6 (9)	7.0 (17)	4.2 (10)	6.1 (14)	3.63 0.06	2.26, 5.82 0.04, 0.08	<0.001 <0.001
OPD	15.3 (11)	6.0 (4)	1.6 (1)	6.4 (4)	1.6 (1)	1.6 (1)	3.3 (2)	1.7 (1)	0.0 (0)			
Quasi Hallucinations												
BPD	41.0 (119)	25.8 (71)	20.1 (54)	11.4 (30)	7.1 (18)	6.8 (17)	6.2 (15)	4.2 (10)	3.5 (8)	3.87 0.06	2.15, 6.97 0.04, 0.10	<0.001 <0.001
OPD	9.7 (7)	6.0 (4)	4.7 (3)	3.2 (2)	3.2 (2)	3.3 (2)	1.7 (1)	1.7 (1)	0.0 (0)			

Table 5: Prevalence of True-Psychotic Thought in Borderline Patients and Axis II Comparison Subject over 16 Years of Prospective Follow-up												
	BL	2 YR FU	4 YR FU	6 YR FU	8 YR FU	10 YR FU	12 YR FU	14 YR FU	16 YR FU	RRR Diagnosis Time	95%CI Diagnosis Time	P-value Diagnosis Time
Any True-psychotic Thought												
BPD	3.8 (11)	1.1 (3)	6.0 (16)	2.3 (6)	1.2 (3)	5.6 (14)	5.7 (14)	6.7 (16)	1.7 (4)	5.18 1.49	1.94, 13.9 0.81, 2.76	0.001 NS
OPD	0.0 (0)	0.0 (0)	1.6 (1)	3.2 (2)	0.0 (0)	0.0 (0)	0.0 (0)	1.7 (1)	0.0 (0)			
True Delusions												
BPD	3.5 (10)	0.7 (2)	4.8 (13)	1.1 (3)	0.4 (1)	3.6 (9)	4.4 (10)	5.9 (14)	0.9 (2)	7.82 1.36	1.92, 31.8 0.63, 2.96	NS NS
OPD	0.0 (0)	0.0 (0)	1.6 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	1.7 (1)	0.0 (0)			
True Hallucinations												
BPD	2.1 (6)	0.4 (1)	3.7 (10)	1.9 (5)	0.8 (2)	4.8 (12)	4.1 (10)	2.1 (5)	1.7 (4)	4.24 1.42	1.34, 13.45 0.68, 2.95	0.014 NS
OPD	0.0 (0)	0.0 (0)	1.6 (1)	3.2 (2)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)			